

Pharmaceutical compositions for treating tumour diseases

The invention relates to the field of tumour therapy.

5 More than 80% of tumours occurring in man are of epithelial origin. The formation of epithelial tumours (carcinomas) is a multi-stage process which is illustrated most clearly in the progression of human colon carcinoma
10 (Powell, et al., 1993) and skin tumours in mice (Wright, et al., 1994). Carcinomas are assumed to start from individual cells or small groups of cells in which mutations have occurred. These cells develop into benign, epithelial hyper- or dysplastic regions. The progression
15 of these hyperplastic regions into a carcinoma *in situ*, which may then acquire invasive and metastatic properties, requires a number of further mutations in the tumour cell. Characteristically these cells acquire the ability to break down their basal membrane proteolytically, to
20 develop from a stationary polarised cell into a non-polarised cell capable of migrating in the tissue, to survive in the bloodstream and form metastases at remote sites (Liotta, et al., 1991; Liotta and Stetler-Stevenson, 1991).

25 Although deep changes in gene expression are involved in the many changes in the architecture and behaviour of malignantly transformed cells, none of these newly acquired properties occurs only in invasive tumour cells.
30 Attachment to the basal membrane, proteolysis thereof and migration through the basal membrane and the underlying mesenchyme are important stages in normal processes, e.g. in the implantation of the trophoblast, in movements of configuration during development of the embryo, in the
35 development of the mammary gland and the reorganisation of epithelia during wound healing (Aznavorian et al., 1993).

For a better understanding of the development and progression of carcinomas it is crucial to understand how the deregulation of these normal processes takes place in cell invasion and metastatisation.

5

Studies in recent years have contributed to an understanding of the molecular mechanisms involved in the modulation of the epithelial phenotype in normal and pathological situations (Reichmann, et al., 1992; Frisch, 1994). Moreover, exogenous polypeptide factors such as Scatter Factor (SF)/Hepatocyte growth factor (HGF) and New-Regulin/HER-Regulin play important roles in the changes in the migration and differentiation properties of epithelial cells (Birchmeier et al., 1993; Hartmann et al., 1994; Soriano et al., 1995). Only recently, Transforming Growth Factor 1 (TGF β 1) was identified as another potent modulator of the phenotype of breast epithelial cells (Miettinen et al., 1994; Zambruno et al., 1995).

20

TGF β 1 belongs to a large super-family of multifunctional polypeptide factors. The TGF β family itself consists of three genes, TGF β 1, TGF β 2 and TGF β 3, which have extremely high homology with one another. In mammals the TGF β -super-family also includes the various TGF β genes as well as the embryonic morphogenes, such as e.g. the family of the activins, "Müllerian Inhibitory Substance", and the bmp family ("Bone Morphogenetic Protein"), which play important roles both in regulating embryo development and in the reorganisation of epithelia (Roberts and Sporn, 1992). TGF β 1 inhibits the growth of many cell types, including epithelial cells, but stimulates the proliferation of various types of mesenchymal cells. In addition, TGF β s induce the synthesis of extracellular matrix proteins, modulate the expression of matrix proteinases and proteinase inhibitors and change the

35

tumour cell, lead to a conversion of epithelial cells into fibroblastoid cells with invasive potential.

2. The autocrine production of TGF β by the converted cells leads to the maintenance of the degenerate, invasive cell status.

3. Interruption of the transmission mediated by the TGF β -receptor signal prevents "epithelial-fibroblastoid conversion" (EFC) and the concomitant invasiveness and may change cells which have already undergone an EFC and are growing in a stably invasive manner back into epitheloid cells which are no longer growing invasively (fibroblastoid-epithelial conversion; FEC).

Within the scope of the present invention, the role of TGF β 1 in the normal development of the mammary glands was investigated with a view to assessing possible side effects of TGF β 1 inhibitors.

Within the scope of the present invention, it was shown, on the one hand, that Ha-Ras-transformed breast epithelial cells (EpRas-cells) undergo a transition (conversion) from the epithelial to the fibroblastoid (or mesenchymal) state in the formation of tumours in mice. This transition is hereinafter referred to as EF-transition or EF-conversion ("Epithelial-Fibroblastoid Cell Conversion", EFC). Such an EF-conversion has also been demonstrated in vitro. For this, EpRas cells were cultivated in type I collagen gels. In the absence of serum these cells developed into three-dimensional, cystic hollow structures, the walls of which consisted of a single-thickness layer (monolayer) of polarised epithelial cells. TGF β 1 caused these same Ras-transformed cells to develop into disorganised strands consisting of spindle-shaped cells with fibroblastoid properties. In non-transformed epithelial cells TGF β 1 was unable to cause such changes. The converted cells were highly invasive both in collagen gels and in chicken heart

- 5 -

- invasion assays. Surprisingly it was found that, once the fibroblastoid cells had undergone the conversion, they themselves produced large amounts of TGF β 1. If this self-produced TGF β 1 was inactivated by a TGF β 1 neutralising antibody, the cells changed back into a polarised, epithelial phenotype. This cell behaviour indicates that the converted fibroblastoid phenotype is maintained by TGF β 1, the TGF β 1 acting through an autocrine loop.
- 10 It was also shown, within the scope of the present invention, that the mechanism observed *in vitro* also applies *in vivo*: tumour cells which had undergone an EF-conversion themselves produced TGF β 1. Moreover, TGF β 1 is capable of triggering and sustaining the invasive
- 15 phenotype of Ha-Ras-transformed breast epithelial cells in experimentally induced tumours.
- Moreover, it was shown within the scope of the present invention that in human tumours of various origins (kidney
- 20 cell carcinoma, breast cancer) there were indications of the occurrence of "Epithelial-Fibroblastoid Cell Conversion" (EFC) (75% of the kidney cell carcinomas investigated and 25-60% of the breast tumours coexpressed the general epithelial marker cytokeratin and the
- 25 mesenchymal marker vimentin). It was also shown that all these tumours themselves produce TGF β 1. This is an indication that the results obtained with the model system used in the present invention also apply to human tumours.
- 30 Fourthly, it has been shown within the scope of the present invention that total inhibition of the signal transmission induced by the TGF β receptor can be achieved using a dominant-negative TGF β -receptor chain II (T β RII-dn). Such expression of T β RII-dn led to the elimination of
- 35 the malignant, invasive phenotype, not only in Ras-transformed mouse-breast epithelial cells, but also in a

number of already mesenchymal, invasively growing carcinoma cell lines in humans and mice and to the complete inhibition of the formation of tumours or metastases obtained by these lines in the experimental animals.

The present invention is thus based on the following findings:

10 Numerous mutations in protooncogenes and tumour suppressor genes participate in carcinogenesis (Vogelstein and Kinzler, 1993). However, little is known about how specific oncogenic mutations are connected with defined changes in the phenotype of the cell and the manner in which these changes then contribute to tumour cell invasion and metastatisation. Within the scope of the present invention it was first demonstrated, using a model system, that the Ras-oncoprotein dramatically changes the cell reaction of breast epithelial cells to TGF β 1 both in collagen gels and in developing tumours. This modified reactivity of the cells causes TGF β 1 to induce an EFC. Once converted, these fibroblastoid cells themselves produced high concentrations of TGF β 1 and thus retained their own mesenchymal and invasive properties.

25 The theoretical validity of this principle could then be demonstrated in a number of unrelated tumour models in humans and mice. In these tumour cells, other oncogenes very probably take on the function of Ha-Ras. It has been shown that in all these cells both TGF β 1 and the interruption of any existing autocrine stimulation by TGF β 1 dramatically influences the tumour cell phenotype: TGF β also leads to an increase in invasive growth in these cells, whereas switching off the TGF β -receptor or the signal transmission pathways activated by it led to re-formation of the EFC, i.e. to a fibroblastoid-epithelial

conversion (FEC) and/or to loss of the invasive, tumour-producing cell phenotype.

5 The experiments carried out within the scope of the
present invention originally started from the observation
that Ras-transformed mouse breast epithelial cells convert
into invasive spindle cells during tumour formation.
Similar spindle cell tumours have been described in the
brain, skin, colon and breast, both in humans and in
10 animal models (Buchmann et al., 1991; Guldberg, 1923;
Sandford et al., 1961; Sonnenberg et al. 1986; Stoler et
al. 1993). The origin of these spindle cell carcinomas is
still unclear, although some researchers believe that
these often highly invasive tumours constitute a separate
15 class of tumour of fibroblastoid origin, whilst other
authors assume that these tumours are of epithelial
origin.

In the model system used within the scope of the present
20 invention the spindle cell tumours first used clearly
originated from the epithelial donor-cells injected into
the animal. Spindle cells originating from the tumour
survived the selection in G418 and expressed cell- and
tissue-specific cytokeratins, confirming their donor cell
25 status and their epithelial origin. Moreover, the tests
carried out showed that the injected epithelial cells and
the converted fibroblastoid tumour cells came from the
same cell clone and re-integration of the retroviral
vector into other sites of the genome could be ruled out
30 as a possible cause of the changes. What was almost more
important was that the fibroblastoid phenotype of the
converted cells was absolutely stable under standard
culture conditions and that the cells changed back to
polarised epithelial cells efficiently after
35 neutralisation of the TGF β 1 activity. This rules out
genetic or epigenetic changes being responsible for the

cell conversion. The most probable explanation for the dramatic change in the phenotype *in vivo* is that an interaction between the Ras-transformed cells and mesenchymal cells surrounding them leads to the conversion of the epithelial cells into fibroblastoid cells. Within the scope of the present invention it has thus been shown that EFC is a mechanism which is relevant to carcinogenesis in certain tumours.

Within the scope of the present invention it has also been shown that TGF β 1 induces EFC both in collagen gels and during tumour development. In the cell model used first, this TGF β 1-induced conversion remarkably requires the co-operation of an activated Ras protein; neither primary breast epithelial cells nor the parental Eph4 cells underwent TGF β 1-induced EFC. From this it can be concluded that EFC is triggered by a synergy of various signal transmission pathways which are activated on the one hand by TGF β 1 and on the other hand by Ha-Ras. This assumption is supported by other findings which indicate that activated Ras proteins have similar effects on cells to members of the TGF β family. This applies e.g. to myogenic differentiation (Payne et al., 1987) and to the formation of the mesoderm (Whitman and Melton, 1992). In the heart muscle, TGF β highly regulates genes associated with the growth of the embryonic heart regulated by haemodynamic loading. These effects are at least partly mimicked by activated Ras (Parker et al., 1990; Thorburn et al., 1993), leading one to suppose that Ras and TGF β can act synergistically, at least in some biological systems.

In connection with this it is interesting that the overexpression of normal and mutated Ras was observed in a considerable number of human carcinomas, including breast cancers (De Bortoli et al., 1985; Hand et al., 1984; Slamon et al., 1984). Furthermore, autocrine production of

ligands as well as overexpression and/or constitutional activation of receptor-tyrosinekinases which occur at the start of signal transmission pathways including c-Ras (e.g. HER-1, HER-2), are frequent changes in breast cancers (Kern et al., 1990; LeJeune et al., 1993). Since TGF β 1 is also abundantly present in many human tumours (Derynck et al., 1985; Keski-Oja et al., 1987; Thompson et al., 1991) it may be concluded, on the basis of the results obtained within the scope of the present invention, that Ras- and TGF β 1-induced signals also act synergistically in human tumours. As demonstrated by the results of the experiments carried out within the scope of the present invention, the TGF β -receptor may also control EFC and invasiveness in tumour cells transformed by oncogenes other than Ras. Another major finding within the scope of the present invention is thus that TGF β co-operates with various oncoproteins, including Ras, and tyrosinekinases, in regulating the plasticity of the polarised epithelial phenotype.

After serum-free (and TGF β -free) cell culture in reconstituted collagen gels, EpRas-cells exhibited a great capacity for organogenesis and a high degree of epithelial polarisation. The Ep-Ras cells, however, predominantly formed widened tubuli as well as alveolar cavities, in contrast to the narrow branching tubuli formed by the parental EpH4-cells or by primary breast epithelial cells. This shows that the Ras-oncoprotein on its own, in the absence of TGF β , is capable of modifying the morphogenetic behaviour of epithelial cells to some extent. In other systems activated Ras has been described as having more powerful effects on epithelial polarity (Eaton and Simons, 1995). Here, transformation with Ras led to disruption of the polar expression of apical proteins, whilst the expression of basolateral marker proteins remained unaffected (Schoenenberger et al., 1994). Since the

- 10 -

experiments described above were carried out in the presence of FCS (which itself contains TGF β 1), however, it is difficult to compare them with the results obtained within the scope of the present invention in which no such
5 obvious polarity defects could be observed. Since exogenous TGF β 1 completely destroyed the cell polarity, it is possible that the partial destruction of the polarity in the abovementioned Ras-transformed cell systems can be put down to the TGF β 1-concentrations present in the serum.
10 Nonetheless the morphogenetic behaviour in EpRas-cells is changed slightly, possibly as the result of increased protease activity.

TGF β completely destroyed the polarity of epithelial cells
15 and caused the cells to become spindle-shaped and invasive. These changes were dependent on the constant presence of TGF β 1, since the spindle cells quickly changed back into polarised epithelial cells when TGF β 1 neutralising antibodies were added. The most important
20 conclusion from these results is that Ras-transformed cells can be switched back and forth by TGF β 1 between a quasi-normal and a highly tumorigenic phenotype. This intense phenotypical plasticity might be characteristic of invasively growing cells in general and might explain why
25 invasive tumour cells often exhibit the migratory properties of fibroblasts. After extravasation these fibroblastoid, migratory cells should be able to develop back into well differentiated secondary tumours in the new environment provided by remotely located tissue (see
30 below). Increased phenotypical plasticity is thus a characteristic of invasive tumour cells.

Another essential finding reached within the scope of the present invention is that EpRas cells have to undergo EFC
35 in order to produce significant amounts of TGF β 1 both in vitro and in vivo. It has been shown that tumour cells are

- 11 -

able to maintain their fibroblastoid phenotype by means of the autocrine production of TGF β 1 and that the autocrine TGF β 1 production and effect on the producing cell (autocrine loop) has to be interrupted in order to make the phenotypical re-conversion of the cells possible. The ability of TGF β 1 to induce EFC and then efficiently maintain the invasive phenotype may also explain why the initially epithelial Ras-transformed cells changed progressively and uniformly into spindle cells during the tumour growth.

Shortly after their injection into mice, polarised Ras-transformed epithelial cells neither expressed nor released significant quantities of TGF β 1. As was established by hybridisation *in situ* and immunohistochemistry, however, the stroma cells surrounding the microtumour expressed the cytokine. These stroma cells could be identified as fibrocyte and endothelial cells, but it must be presumed that other cell types, such as macrophages and lymphocytes, were probably also present; all these cell types are known to produce and release TGF β 1. The most probable conclusion is that the effects of TGF β 1 are regulated primarily at the level of their proteolytic activation. The primary regulation of TGF β is carried out by factors which control the processing of the latent into the biologically active molecule. However, virtually nothing is known about the TGF β activation *in vivo*. The protease plasmin can activate two cell types of latent TGF β 1 in co-culture systems, but only if two different cell types are in direct contact or close together (Antonelli-Orlidge et al., 1989; Sato et al., 1990). This close contact of different cell types should take place in the system used within the scope of the present invention after encapsulation of the tumours by the stroma and to an even greater extent if donor-tumour cells are mixed with stroma cells of the receiver

animal during the tumour development (Fig. 2B). Furthermore, thrombospondin (TSP), an extracellular matrix protein, activates latent TGF β . In this case the activation takes place in the soluble phase and requires
5 no proteolytic activity (Schultz-Cherry et al., 1994). In fact, the role of thrombospondin in supporting the development of cancer and the increased thrombospondin concentrations in malignant breast cancers have been briefly reported (Castle et al. 1991; Wong et al., 1992).
10 It has thus been shown within the scope of the present invention that the autocrine production of TGF β 1, in cooperation with the oncoprotein Ha-Ras, maintains the fibroblastoid phenotype.

15 The findings and conclusions reached within the scope of the present invention led to the hypothetical model shown in Fig. 9. It is postulated that the TGF β 1 which is relevant to the tumour formation is produced primarily by infiltrating cells of the tumour stroma, such as
20 fibrocytes, endothelial cells, lymphocytes and macrophages. The interaction of the tumour cells with the different cell types of the tumour stroma should trigger the efficient production and/or activation of TGF β 1. This should in turn cause the epithelial tumour cells to change
25 into the fibroblastoid and invasive phenotype. These fibroblastoid cells themselves then start to produce TGF β 1 which acts on them in an autocrine loop and thus both maintains the fibroblastoid phenotype and also makes it easier to recruit other epithelial cells to the EFC.

30 Further mutations or selective mechanisms should cause some of these invasively growing cells to migrate into blood vessels and out again and finally to form secondary tumours at remote sites. This model conforms to findings which show that increased TGF β 1 expression is also
35 involved in the progression to malignancy in a murine prostate cancer model (Thompson et al., 1992; 1993).

- 13 -

The findings described hitherto were obtained in a combined *in vitro/in vivo* model system using Ras-transformed mouse-breast epithelial cells. Within the scope of other tests, crucial aspects of this model (EFC, TGF β 1 production in the tumour) were detected in a large number of primary human carcinomas of the kidneys and breast. Thus, in the majority of all the kidney cell carcinomas investigated as well as in a percentage of the breast tumours investigated dependent on the degree of malignity, the occurrence of an EFC is demonstrated by the coexpression of cytokeratin (general epithelial marker) and vimentin (mesenchymal marker). Moreover, in all the tumours investigated, the production of TGF β 1 by the tumour cells themselves has been demonstrated both at the protein level by histochemical staining with anti-TGF β -antibodies and also at the mRNA level by *in situ* hybridisation and RT-PCR.

By means of another series of tests carried out within the scope of the present invention it has been shown that the TGF β -receptor generally assumes a central position in the regulation of EMT and invasive tumour cell growth. Not only in Ha-Ras transformed breast epithelial cells, but in a number of other tumours which originate from other epithelial types and wherein it is not known which oncogenes take over the function of Ha-Ras, the TGF β -receptor has been identified as the crucial regulator of epithelial plasticity as well as of the invasive growth of the tumour cells. Thus, it has been possible to completely inhibit the invasive growth of two human carcinoma cell lines (kidney carcinoma line MZ 1795, nasopharyngeal carcinoma line KB) (presumably caused by secreted TGF β 1) in collagen gels by means of a neutralising anti-TGF β 1-antibody.

- 14 -

The proof of the above hypothesis was finally provided by means of a dominant-negative TGF β receptor (T β RII-dn). This T β RII-dn constitutes a so-called "kinase-dead" mutant of the receptor chain II which binds to endogenous
5 receptors of type I, but cannot phosphorylate them. In this way all the T β RII-dn-bound TGF β -receptor chains of type I are inactivated because they cannot activate any signal transmission even after the binding of the ligand (TGF β 1) since the phosphorylation by receptor chain II
10 required for this is absent. If a dominant-negative TGF β -receptor of this kind is overexpressed in tumour cells, the entire signal transmission proceeding from the TGF β -receptor can be inhibited in these cells. The expression of T β RII is thus suitable for simulating the activity of
15 inhibiting TGF β or inhibiting the signal transmission pathway triggered by the activation of the TGF β -receptor.

At first T β RII-dn was overexpressed in Ha-Ras-transformed mouse-breast epithelial cells (EpRas). All the clones
20 obtained exhibited greatly delayed tumour growth in nude mice. Moreover, the cells isolated from such tumours had an epithelial phenotype and expressed epithelial markers (E-Cadherin, ZO-1) but no mesenchymal markers (vimentin). This shows that the expression of a T β RII-dn inhibited EFC
25 during tumour formation.

After obtaining these results it was useful to check whether switching off the signal transmission of the TGF β -receptor also works in tumour cells which have already
30 undergone EFC and thus have a stable mesenchymal and invasive phenotype. The colon carcinoma line CT26 in the mouse was selected as an example of such a cell line. This tumour cell line has a very marked tendency to form lung metastases rapidly after subcutaneous injection in mice,
35 so that the animals die from the lung metastases even after the primary tumour has been surgically removed in

- 15 -

good time. This cell exhibits mesenchymal morphology, grows into disordered chains and strings of spindle-shaped cells in the collagen gel and expresses no epithelial markers apart from basal cytokeratins. Instead the cells
5 have a high vimentin expression. If the dominant-negative TGF β -receptor (T β RII-dn) is overexpressed in these cells, the cells form smaller or larger compact clumps in the collagen gel and grow on plastic as epitheloid cells which form hemicysts (domes) and express large amounts of E-
10 cadherin and ZO-1. The cells were thus obviously changed back, by the T β RII-dn, into cells with an epithelial phenotype (fibroblastoid-epithelial conversion, FEC).

A corresponding activity of the dominant-negative TGF β -
15 receptor (T β RII-dn) was also observed *in vivo*. When different, T β RII-dn expressing clones of CT26 cells were injected into mice, the tumour formation was delayed by different amounts depending on the clone. In many clones tumour formation only occurred after 6-8 weeks, as opposed
20 to 1-2 weeks in the case of animals injected with control CT26 cells without T β RII-dn. However, the activity of the T β RII-dn was even more dramatic when the primary tumours were removed from the mice at a certain size and the formation of metastases was expected. In this experiment
25 metastases did not develop in any of the mice injected with T β RII-dn expressing CT26 cells (even after more than 18 weeks) whereas the control animals died of lung metastases within 2-4 weeks after excision of the tumour.

30 The decisive conclusion from these experiments for the present invention is that inhibiting the signal transmission mediated by the TGF β -receptor can not only prevent the occurrence of an EFC and the resulting acquisition of invasive properties, but also change any
35 existing, invasively growing tumour cells back into a benign state in which they are no longer invasive.

- To sum up, the findings obtained within the scope of the present invention indicate that the increased sensitivity and altered reactivity of the cells, compared with the ability of TGF β to modify the epithelial phenotype, generally represents a characteristic of epithelial tumour cells. This altered reactivity can be brought about by Ras-(onco)proteins, but also by tyrosinekinases which activate Ras, as well as by other as yet unknown oncoproteins. This altered oncogene-induced mode of reaction to normal environmental signals, such as e.g. those induced by TGF β 1, should lead to altered gene expression in the tumour cell and also incorrect transmission or interpretation of signals between tumour and stroma cells. This abnormal "crosstalk" between tumour cells and their immediate environment would appear to be the driving force for what is commonly known as tumour progression.
- Furthermore, the results of the experiments show that activated Ras, overexpressing receptor-tyrosinekinases which activate the Ras signal transmission pathway as well as other, possibly unknown oncogenes co-operate both in the normal development and also in the carcinogenesis with the TGF β 1-receptor. This would appear to involve processes such as the induction/activation of stromal TGF β 1 by the interaction of epithelial and mesenchymal cells as well as EF conversion, induced and maintained by TGF β 1.
- The main difference between normal and oncogene-transformed tumour cells should therefore be as follows: TGF β 1 has a physiological, strictly regulated function during the morphogenesis of normal cells. In the tumour cell the transformation by oncogenes causes degeneration of the function of TGF β 1, i.e. constitutive, highly abnormal morphogenetic changes are triggered in the cells.

- 17 -

The present invention thus relates to a pharmaceutical composition containing as active compound a substance which inhibits the activity of TGF β on tumour cells of epithelial origin, for treating epithelial, invasive tumour diseases which are characterised by a reversible transition of the cells from an epithelial, non-invasive state into an invasive state.

10 In one embodiment of the invention the pharmaceutical composition also contains substances which inhibit the expression of oncogenic Ras and/or the overexpression of normal Ras or the activity of Ras-activating receptor tyrosinekinases in the cells.

15 In epithelial invasive tumour diseases the tumour cells have an increased phenotypical plasticity, i.e. they are able to undergo transitions from the epithelial, non-invasive state to the fibroblastoid, invasive state (EF conversion) and *vice versa* (FE conversion).

20 The substance which inhibits the activity of TGF β on the cells or the signal transduction mediated by activation of the TGF β -receptor is hereinafter referred to as "TGF β inhibitor".

TGF β s, like the other members of the TGF β super-family of multifunctional polypeptide factors such as e.g. activins, Bone Morphogenetic Proteins (bmp's), etc., exert their effect by binding to specific cell surface receptors. The type I and type II TGF β receptors form heterodimeric complexes after binding of the ligand, thereby initiating the signal transmission. The type II receptors, which are assigned to the group of the receptor serine/threonine-kinases in terms of their activity, bind the ligands, but require association with the type I receptors which

- 18 -

constitute serine-threonine-kinases in order to be able to pass on the signal obtained from the ligand. Whereas the type II receptors are responsible for the ligand specificity, the functionally different type I receptors heterodimerise with several type II receptors. In this ligand-induced heterodimerisation the type II-receptor chains phosphorylate the type I receptors on serine/threonine groups and thereby activate them. This cooperation of the type II-receptor with a particular type I-receptor causes activation of specific signal transmission pathways and as a result leads to a transcriptional response to the signals transmitted to the cell by the ligands.

The activity of a TGF β inhibitor is based on the fact that it blocks the cell response triggered by the receptor activation, i.e. it prevents the TGF β -receptor system from being activated and hence the cell signal transmission pathway from being actuated.

Since it is the type I receptors which are responsible for the specific transcriptional response which eventually produces the fibroblastoid phenotype after the binding of the ligands to the type II receptor, and to the type I receptor, the type I receptor represents one of the target molecules for the TGF β inhibitor. Because of the need for phosphorylation of the type I receptor by the type II receptor (and on the basis of the results obtained with dominant negative type II receptors whose serinekinase activity has been destroyed by mutation, the type II receptor is also a possible target molecule for inhibitors.

Other mechanisms for the activity of a TGF β inhibitor are thus based on preventing the interaction between the ligand TGF β and the type II receptor, preventing the

- 19 -

signal transmitted from the type II receptor to the type I receptor which brings about the activation of the type I receptor. Finally, blocking the binding of TGF β to the type I receptor, inhibiting the activity of the type I receptor or inhibiting an effector molecule of the signal transmission pathway activated by the type I receptor are all possible methods of attacking inhibitors.

Examples of inhibitors are antibodies which neutralise the TGF β , particularly monoclonal antibodies, TGF β antisense-RNA molecules (Fakhrai et al., 1996) or dominant-negative TGF β receptors of type I or II.

The invention relates, according to a further aspect, to screening processes for identifying pharmacologically active substances for treating epithelial, invasive tumour diseases which are characterised by a reversible transition of the cells from an epithelial, non-invasive state into an invasive state.

One method of finding suitable, particularly low-molecular, inhibitors, comprises determining, in a first step, which of the type I receptors is responsible for the transition from the epithelial to the fibroblastoid state of the cells. To do this, the EpRas-cell line used within the scope of the present invention (or one of the other cell lines used which are capable of bringing about the EF conversion or have already undergone one), is interrogated to see which TGF β -type I/II receptor it expresses. This interrogation may be carried out by RT-PCR ("Reverse Transcriptase Polymerase Chain Reaction") using oligonucleotides, derived from known TGF β -type I or type II receptors, as PCR primers in order to amplify the relevant receptor-DNA from EpRas-DNA and thus identify the TGF β -type I or type II receptor expressed in these cells. The experiments described in the Examples with a dominant-

- 20 -

negative mutant of the human type II receptor T β R-II (only this chain occurs in all the known receptors for TGF β 1,2,3; Wrana et al., 1992, Wrana et al., 1994) confirm that this TGF β -type II receptor subtype (as such) is
5 necessary, directly or indirectly, for the signal transmission which leads to the EF conversion. This TGF β -type II receptor subtype thus constitutes one of the target molecules for the TGF β inhibitor. This satisfies an essential precondition for establishing a cellular or
10 biochemical screening assay which can be used to screen specifically for substances which inhibit this target molecule.

Next, an investigation is carried out in a cell which is
15 undergoing the EF conversion or in which the EF conversion can be reversed by inhibiting the TGF β receptor signal transmission to determine which of the processes taking place in the cell by EF conversion or reversal thereof is most suitable for establishing a screening assay.
20 Appropriately the EpRas cells used in the Examples or the CT26 cells which are also well characterised within the scope of the present invention may be used.

The effects to be expected can be divided into two groups:
25 the first group includes the effects of TGF β on normal mesenchymal and epithelial cells, e.g. in wound healing, described in the literature. The second group of changes are those which occur in particular with the activity of TGF β on transformed cells (such as e.g. in the experiments
30 described within the scope of the present invention). Whereas the TGF β effects of the first group can be adduced for a primary HTS screen ("High Throughput Screen"), any inhibitor candidates found must be tested without fail for their inhibiting activity on the TGF β effects of the
35 second group.

- 21 -

The TGF β effects of the first type include i) the induction of extracellular matrix proteins, such as fibronectin, laminin, elastin; ii) the induction of the protease inhibitor PAI (Plasminogen Activator Inhibitor) and hence the inhibition of cell protease activity, and iii) an inhibition of cell growth and induction of programmed cell death (apoptosis) in certain cell types. These include in particular normal epithelial cells as well as only slightly degenerate, essentially still epitheloid tumour cell lines. The induction of PAI-1 expression as well as the TGF β -induced apoptosis are possible procedures which may be used to design a cell assay system for screening substances which may act as TGF β receptor inhibitors. The effect chosen is used directly as a system for demonstrating the inhibiting activity of the substance.

In order to find out whether the EF conversion in the EprAs-cell line used within the scope of the present invention triggered by the activation of the TGF β receptor system is transmitted via the same type I/type II receptors as the induction of PAI (or another molecule regulated by TGF β) in untransformed cells, e.g. in the normal starting cell line Eph4 (also used within the scope of this invention), it is possible to check e.g. whether the induction of PAI (or another molecule) or the growth inhibition which is very marked in this cell line is blocked by a dominant-negative mutant of the same type I or II receptor which also blocks the EF conversion. In the case of the CT26 cells which overexpress the dominant-negative type II receptor (T β R β II-dn), it has been shown that, in the T β R β II-dn expressing CT26 clones which reverted to epithelial cells, activation of a PAI-1 promoter-controlled reporter gene was completely inhibited by TGF β -1. The extent of the PAI-I inhibition of the inhibition of reporter gene expression by PAI-I correlated

- 22 -

directly with the ability of the different clones to form tumours in the animal. Moreover, a special CT26 clone, which had recovered complete TGF β -1-inducibility of the PAI-1 promoter-reporter gene construct after lengthy
5 passage *in vitro* (presumably by repressing the T β RII-dn expression) also regained the ability to form metastasising tumours in the mouse.

The confirmation of the correlation between EFC, tumour
10 formation and TGF β receptor type II function using the T β RII-dn experiments provides the prerequisite for a screening assay based on a PAI-1 reporter gene test cell. This test cell, which is a human or animal cell, is stably transformed with a plasmid, in which a reporter gene, e.g.
15 the luciferase gene, is under the control of the regulatory sequence of the PAI gene (or a gene which codes for another molecule regulated by TGF β , e.g. for an extracellular matrix protein). The test cell is also transformed with the human type I or type II receptor,
20 which was shown, after further tests, to be most efficient both at triggering the EF conversion and also at inducing PAI or another molecule regulated by TGF β . The human TGF β type II receptor used for the construction of the T β RII-dn is one of the possible target molecules for a
25 TGF β inhibitor. The control cell used is expediently a parallel-cell clone in which the PAI-1-promoter controlled reporter gene is activated by another receptor not related to the TGF β receptor (e.g. members of the FGF (fibroblast growth factor) receptor-tyrosinekinase family).

30 If a substance which wholly or partially inhibits the TGF β -induced reporter gene expression is found in a screening assay of this kind, it can be concluded that either the selected ligand-activated type I/type II
35 receptor or the signal transmission mediated by this receptor is blocked by this substance. The same substance

should not have any influence on the slight basal reporter gene expression in the control cell in which the reporter gene has been activated not by TGF β , but by FGF. The test systems in which the reporter gene activation is measured
5 can be used in robotised High Throughput Screen (HTS) processes.

A second possible way of measuring the blocking of the TGF β receptor function by test substances can easily be
10 measured by the removal of the growth inhibition and apoptosis brought about by TGF β . Since TGF β efficiently induces apoptosis in normal Eph4 cells under certain conditions, effective inhibitors of the TGF β receptor should act as survival or growth stimulating factors.
15 Eph4 cells in which another apoptosis-inducing receptor has been expressed may be used as control cells. The Fas receptor, which efficiently induces apoptosis in virtually all cell types after the binding of a special Fas ligand, is particularly suitable. The removal of an apoptotic
20 effect by effective TGF β receptor inhibitors has the advantage that it can easily be measured in commercially obtainable test systems (e.g. in the MTS assay which detects the number of live, metabolically active cells), and that toxic substances (which cause rather than prevent
25 cell death) can easily be identified as such. Thus, this test system is also suitable for HTS primary screens.

Another possible cell assay system with which substances can be tested for their inhibiting activity on the EF
30 conversion triggered by activation of the TGF β receptor system, is based on the expression of proteins which are characteristic of the fibroblastoid cell type after EF conversion and are thus an indicator of the occurrence of EF conversion. One example of this is vimentin (Reichmann
35 et al, 1992): it has been shown within the scope of the present invention that expression thereof goes hand in

- hand with the EF conversion triggered by cooperation of Ras and TGF β . Other examples of other markers of the fibroblastoid phenotype are the loss of the expression of E-Cadherin mRNA as well as the *de-novo* expression of
- 5 fibronectin and diverse proteases (UPA, TPA, Reichmann et al, 1992). A suitable test cell transformed by Ras or another oncogene is transformed with a plasmid in which a reporter gene is under the control of the vimentin gene promoter or of promoters of one of the other fibroblastoid
- 10 marker genes mentioned. The modulation of the reporter gene expression by a test substance should then correlate with the modulation of the EC conversion brought about by the same inhibitors.
- 15 Another possible way of finding substances which inhibit the activation of the TGF β receptor system uses the expression of TGF β itself as a detection system. This assay principle is based on the finding reached within the scope of the present invention that the activation of the
- 20 TGF β receptor system in oncogene expressing cells by the ligand TGF β causes the autocrine production of TGF β which acts on the cells in an autocrine loop. In an assay of this kind, which can detect both the activation of the TGF β receptor system and also the induction of the
- 25 autocrine TGF β loop brought about by the expression of Ras (the activity of substances which inhibit TGF β expression, in a test of this kind, do this on the basis of their effect on the activation of the TGF β receptor system and their effect on Ras), the cells contain a reporter gene
- 30 construct which is under the control of the TGF β gene promoter (Kim et al, 1989).
- Biochemical assays in which TGF β inhibitors are identified, the activity of which is based on the fact
- 35 that they inhibit the TGF β signal transmission pathway, may be carried out as follows, for example: in an assay

- 25 -

format the autophosphorylation of the TGF β receptor type II or the cytoplasmic domain thereof which contains the kinase domain is measured *in vitro* on serine or threonine groups, in the presence and in the absence of test substances (potential TGF β inhibitors), a kinase assay of this kind being carried out using methods known from the literature, e.g. as described by Lin et al., 1992, or Braunwalder et al., 1996, and using a receptor (or a domain thereof) prepared by recombinant methods, e.g. in E. coli. In an alternative assay format, the ability of the TGF β receptor type II to phosphorylate the TGF β receptor type I or its so-called GS domain (Wrana et al., 1994), is measured, again according to the known principle of kinase assays, in the presence and absence of potential inhibitor. The modification of an assay of this kind for a High Throughput Format can be carried out using commercially available technologies such as filter plates, FLASH plates (Amersham) or SPA (Scintillation Proximity assay)-Beads (Amersham).

In one of the test systems described, inhibitors of the TGF β receptor found in the primary screen are expediently tested for their specificity in secondary screens. This can be done particularly by direct inhibition of the TGF β -dependent EF conversion of EpRas cells in collagen gels. Another possibility is the incubation of converted EpRas cells (e.g. from mouse tumours) plated out at low density on plastic dishes with the inhibitor of the TGF β receptor found. Effective substances should trigger the conversion of fibroblastoid into epithelial cells even in the presence of TGF β . The same substances should cause re-epithelialisation (FE conversion) in CT26 cells. Finally, particularly suitable active substances in mice which were injected with CT26 cells can be tested to see whether they slow down the growth of the primary tumour or metastasisation after excision of the primary tumour.

The substance which inhibits the expression or the function of oncogenic Ras and/or the overexpression of normal Ras (or the consequences of this overexpression) and/or the activation of normal Ras by receptor tyrosinekinases in the cells is hereinafter referred to as "Ras inhibitor".

Ras inhibitors for the purposes of the present invention either inhibit Ras directly, by inhibiting the activation/function of Ras itself or by inhibiting the activation/function of a Ras-effector molecule which acts below Ras in the Ras signal transmission pathway. Examples are inhibitors of Raf, such as Raf antisense-oligonucleotides (Monia et al., 1996). For cases where the activation of Ras cannot be put down to a change in Ras itself, but it due to the constitutive activation of receptor-tyrosinekinases acting above Ras, inhibition of Ras-activation can also be brought about by inhibiting these receptors. Examples of receptors of this kind are the receptor-tyrosinekinases EGF receptor ("Epidermal Growth Factor Receptor") and homologous receptors such as HER-2, HER-3 or HER-4. Examples of chemical compounds which inhibit the EGF receptor can be found in WO 96/07657. Known Ras inhibitors are monoclonal antibodies (Furth et al., 1982), dominant-negative mutants (Stacey et al., 1991; Quilliam et al., 1994) and antisense-RNA. Examples of low-molecular Ras inhibitors are inhibitors of Ras-Farnesyl transferases (Kohl et al., 1993; Kohl et al., 1994; Kohl et al., 1995).

In order to screen for other low-molecular Ras inhibitors, genes coding for mutations of the Ras proteins H-Ras, K-Ras or N-Ras, which lead to constitutive activation of Ras, are introduced into mammalian cells, e.g. by means of retroviral vectors, and the selective cytotoxic activity

- 27 -

of test substances on the ras-transformed cells is determined. A suitable method of identifying ras inhibitors is described e.g. in der EP-A 604 181.

- 5 Examples of Ras-transformed cell lines which may be used as test cells for the identification of Ras inhibitors, have also been described by Andrejauskas and Moroni, 1989, as well as by Jenkins et al., 1993.
- 10 Ras inhibitors can also be identified with an assay based on the EpRas-cell line used within the scope of the present invention. For this, the cells contain a reporter gene construct in which the reporter gene is under the control of the regulatory sequence of the TGF β gene. First
- 15 of all TGF β is applied to the cells in order to bring about the EF conversion. Then the cells are treated with the test substances. Test substances which are capable of inhibiting the activity of the reporter gene can be assumed to be Ras inhibitors. This can then be confirmed
- 20 in secondary screens in which the substances are investigated to see whether they can inhibit the TGF β - induced EF conversion of EpRas cells in collagen gels or reverse the EFC which has already taken place.
- 25 The pharmaceutical compositions according to the invention can be used, firstly, to prevent the cells from changing into the fibroblastoid state and becoming invasive, thus preventing or reducing their tumorigenicity. Secondly, the pharmaceutical compositions according to the invention can
- 30 also be used to bring about the conversion of existing fibroblastoid and invasively growing tumour cells into non-malignant or less malignant epithelial cells.
- The pharmaceutical composition according to the invention
- 35 may be used on the one hand to prevent the transformation of the cells from the epithelial, non-invasive state into

- 28 -

a fibroblastoid, invasive state. One example of this is its administration after surgical removal of a primary tumour to prevent any tumour cells present from becoming invasive and producing further tumours by metastatisation.

5 Moreover the pharmaceutical composition according to the invention can also slow down tumour growth by the same mechanism, as has been shown with the aid of the T β RII-dn expressing CT26 cells.

10 The pharmaceutical composition according to the invention may, on the other hand, be used to reverse an EF conversion of the cells which has already taken place. Once the conversion has taken place, TGF β maintains the fibroblastoid state by means of an autocrine loop. The
15 administration of a TGF β inhibitor on its own in this case switches off the autocrine loop and thus reverses the fibroblastoid, invasive state of the cell into the normal, epithelial state. However, this reversal is temporary, and there is no fundamental change to the transformed state of
20 the cell brought about by Ras or other oncogenes. This means that when the TGF β inhibitor is removed the EF conversion could start up again. If, on the other hand, an oncogene inhibitor, e.g. a Ras inhibitor or an HER-1/2 inhibitor, is administered, possibly in addition to the
25 TGF β inhibitor, the transformed state of the cell is cancelled, the cell behaves like a normal epithelial cell and reacts correspondingly normally to TGF β , i.e. the effect of TGF β on the cell cannot bring about EF conversion and even leads to growth inhibition of the
30 tumour cell.

The conjecture that TGF β (receptor) inhibitors could cause slowing down or even inhibition of tumour growth is supported by the following state of affairs: most tumours
35 constantly produce TGF β (see below) which is released into the environment and has an immunosuppressant effect there,

i.e. inhibits the function of cytotoxic T-lymphocytes and other cells of the immune system. If the TGF β receptor inhibitor causes the transformation of invasive tumour cells into non-invasive, more epithelial cells, these
5 should switch off the secretion of TGF β and thus be more easily attacked and lysed by cytotoxic T cells.

In order to achieve optimal activity, the pharmaceutical composition according to the invention preferably contains
10 a combination of TGF β inhibitor and Ras inhibitor.

In the transition of epithelial cells into the fibroblastoid state, fibroblastoid marker proteins, e.g. vimentin, are expressed more intensely. The increase in
15 the expression of these markers (see below) is thus one of the diagnostic parameters for tumour diseases which can be treated using the pharmaceutical composition according to the invention.

20 These tumour diseases include adenocarcinomas of the breast (Heatley et al., 1993), kidney cell carcinomas (Beham et al., 1992), carcinosarcomas of the breast (Wargotz and Norris, 1989), carcinosarcomas of the oesophagus (Guarino et al., 1993) or of the female genital
25 tract (de Brito et al., 1993), epitheloid sarcomas as well as spindle cell carcinomas of various locations, e.g. lung carcinomas with spindle cell components (Matsui et al., 1992) or spindle cell carcinomas of the gall bladder (Nishihara et al., 1993).

30 The pharmaceutical composition according to the invention is preferably used to treat breast tumours and kidney cell carcinomas.

35 The pharmaceutical compositions according to the invention are administered to humans in doses of 0.01 to 100 mg/kg

body weight, preferably 0.1 to 15 mg. Apart from the active compounds the pharmaceutical composition contains the usual inert carriers and excipients. The skilled person will find methods of formulating pharmaceutical preparations in the relevant textbooks, such as Remington's Pharmaceutical Sciences, 1980.

10

Figure Summary

- Fig. 1: conversion of EpRas cells into fibroblastoid cells during tumour formation in mice
- Fig. 2: epithelial/mesenchymal conversion (EFC) during tumour development: time scale and behaviour of donor and receiver cells
- Fig. 3: organogenesis and epithelial polarity are destroyed by serum or TGF β 1
- Fig. 4: TGF β 1 destroys the cell polarity in Ras-transformed breast epithelial cells
- Fig. 5: fibroblastoid EpRas cells are highly invasive in the chicken embryo heart invasion assay
- Fig. 6: TGF β 1 maintains the fibroblastoid phenotype of converted EpRas cells via an autocrine loop
- Fig. 7: converted EpRas cells produce high concentrations of TGF β 1
- Fig. 8: TGF β 1 triggers the transition from the epithelial to the fibroblastoid state as well as the invasiveness of the cells in experimentally induced tumours
- Fig. 9: model for the activity of TGF β 1 in tumour development
- Fig. 10: TGF β 1 induces *in vitro* morphogenesis and apoptosis in normal mammary gland epithelial cells

- Fig. 11: *In vivo* expression of TGF β 1 during the formation of the normal breast
- Fig. 12: *In vivo* expression of TGF β 1 during the breakdown of the fully developed mammary gland after ablation
- 5 Fig. 13. inhibition of the invasivity of human tumour cells by TGF β -neutralising antibodies
- Fig. 14 expression of a dominant-negative TGF β -receptor (T β RII-dn) in Ras-transformed breast epithelial cells inhibits EFC and tumour growth
- 10 Fig. 15 expression of T β RII-dn in mouse-colon carcinoma cells (CT26) prevents growth in collagen gels and invasivity
- 15 Fig. 16 expression of T β RII-dn in CT26 cells inhibits metastatisation *in vivo*
- Fig. 17 T β RII-dn expressing CT26 cells are incapable of forming metastases in the lung even after intravenous injection
- 20 Fig. 18 expression of a PAI-1-promoter-reporter construct in CT26- and CT26-T β RII-dn cells]

25 In the following Examples, the following materials and methods were used unless otherwise stated:

a) Cell culture

- 30 EpRas cells were prepared by infecting the parental breast epithelial cell line EpH4 (a subclone of the spontaneously immortalised breast epithelial cell line Ep1 (Reichmann et. al, 1992) selected for the strong impression of a polarised phenotype) with a helper-free, v-Ha-Ras
- 35 expressing retroviral vector (Redmond et al., 1988). The selection and expansion of the polarised epithelial clones

- was carried out as described by Reichmann et al., 1992. For this the cells were cultivated on plastic dishes in growth medium (Dulbecco's modified Eagles medium; (DMEM), containing 10% FCS (Boehringer Mannheim) and 20 mM HEPES, and subcultivated in a ratio of 1:3 twice a week. For the induction of hemicyst(dome) formation EpRas cells and cells of the parental line EpH4 were cultivated at high density for one week without subcultivation.
- 10 The human tumour cell lines MZ 1795 (kidney carcinoma; Seliger et al. 1996) and KB (nasopharyngeal carcinoma ATCC CCL17; Derynck et al., 1985) were obtained from the ATTC. They were cultivated in the same medium as the mouse EpH4 cells.
- 15 The mouse colon carcinoma line CT26, the establishing of which was described by Brattain et al., 1980, was also cultivated in the same medium as the mouse-EpH4 cells.
- 20 In order to express the human, dominant negative TGF β receptor type II (T β RII-dn, Wrana et al., 1992) in mouse-EpH4 and CT26 cells, the corresponding cDNA (Wrana et al., 1992) was inserted in the helper-free retrovirus vector pbabe-puro (Morgenstern and Land, 1990). The retrovirus-DNA was transfected into BOSC 23 packaging cells (Pear et al, 1993) and the virus-producing, Mitomycin-C treated BOSC 23 cells cocultivated with EpH4 or CT26 cells. Infected clones were selected with G418 and expanded in Dulbecco's modified Eagles medium (DMEM), containing 20% FCS (Boehringer Mannheim) and 20 mM HEPES. The expression of the T β RII-dn protein was detected in the Western Blot using a haemagglutinin (HA) epitope present in the construct.
- 30
- 35 b) Growth of organotypical cell structures in collagen gels

- 33 -

Semiconfluent cultures of the cells to be analysed were trypsinised and adjusted to a final concentration of 4×10^4 cells per ml with ice-cold growth medium. Equal
5 volumes of the cell suspension and an acidified solution of rat's tail collagen type 1 (Sigma) were mixed at 4°C , applied to 35mm tissue culture dishes and incubated for 30 min at 37°C , in order to allow the solution to set to a gel. To allow the cells to form organotypical structures,
10 the collagen gels were covered with a serum-free medium (MEGM; Promocel) containing bovine pituitary extract (BPE), recombinant epidermal growth factor (EGF), hydrocortisone and insulin (in the concentrations recommended by the manufacturer). Where stated, 5 or 10% FCS or 5ng/ml
15 recombinant TGF β 1 (literature) were added. The medium covering the collagen gels was changed every two days. In order to neutralise the TGF β 1 produced by the cells or cell structures in the collagen gels a monoclonal antibody against TGF β 1 (Genzyme) or control antibody is used in
20 concentrations of up to $50\mu\text{g/ml}$.

c) Tumour induction in mice and re-isolation of cells from tumours or collagen gels

25 Confluent EpRas or Eph4 cells were trypsinised and counted. Then 10^5 cells, suspended in 0.1ml PBS, were injected subcutaneously or into the mammary gland of 5 week old BALB/c mice or nude mice. The mice were killed after different lengths of time (between 3 and 28 days)
30 and the tumours (or tissue zones which the injected cells contained) were excised. For the subsequent histological analysis the tissue was immediately flash-frozen in liquid nitrogen. To isolate the tumour cells for further growth in the tissue culture, the tissue was cut into small
35 pieces under sterile conditions using two opposing scalpels and digested with 2mg/ml collagenase type 1

- (Sigma) for 1 hour at 37°C. In order to remove the remaining host cells which lacked the retroviral neomycin or hygromycin resistance markers of the donor cells, the cells obtained from the tumours were grown for the first 5 days in the presence of G418 or hygromycin. The collagen gels were digested with collagenase in a similar manner, in order to isolate the cells for the subsequent tissue culture.
- 10 For tumour induction with CT26 cells or CT26-T β RII-dn cells, 1×10^6 cells per animal were injected subcutaneously into nude mice under the skin of the back. The size of the detectable tumours was determined every 3 days and the animals were killed when the tumours exceeded a size which was tolerable for the wellbeing of the animals.
- 15 In another experiment 1×10^6 cells per animal were injected into syngenic mice (Balb/C). After the primary tumour reached a size of 4 cm³ the tumours were surgically removed so that there was no tumour tissue remaining at the site of the operation. The mice were then further monitored and after their death the presence of lung metastases was investigated.
- 20 In a last experiment to demonstrate the ability of the CT26- and CT26-T β RII-dn cells to colonise the lung from the circulation, 5,000 and 50,000 cells of both types were injected i.v. into the caudal vein of syngenic Balb/C mice. After their death the mice were then examined as before for the presence of lung metastases.
- 25
- 30 d) Antibodies
- Rabbit antisera against cytokeratin have been described by Reichmann et al., 1992. Rabbit antiserum and monoclonal rat antibodies against E-Cadherin were prepared as described by Kemler, 1993 and the relevant literature cited therein (Kemler, 1993). Rabbit antiserum which
- 35

recognises neomycin phosphotransferase was prepared by bacterially expressing neomycin phosphotransferase, purifying it and injecting it into rabbits. After a reasonable time rabbit serum was obtained and

- 5 used neat for immune staining. The monoclonal mouse antibody against vimentin V3B (Boehringer Mannheim), the monoclonal rat antibody against ZO-1 (Chemicon), the monoclonal mouse antibody against TGF β 1-3 (Genzyme), the TGF β 2,3-antibody (Genzyme), the polyclonal antiserum
10 against activated TGF β (Promega), the TGF β -neutralising polyclonal rabbit antibody (R&D) as well as the monoclonal TGF β -antibody (Genzyme) were obtained commercially.

e) Sections and immunofluorescence

- 15 In order to obtain the optimum biological activity from RNA and proteins in the excised tissues and collagen gels, the tumour material and the cell structures containing collagen type I gels were flash-frozen immediately after
20 isolation in liquid nitrogen. Before freezing the collagen gels were soaked for 2 min in medium containing 5% DMSO in order to prevent cell damage as a result of the formation of ice crystals. Cells grown on plastic or frozen sections prepared from tumours or collagen gels were fixed and made
25 pervious for 15 min at -20°C using acetone/methanol, mixed in the ratio 1/1, air-dried and stored at 4°C. The incubation with the first antibody was carried out for 1 h at 37°C in PBS which normally contained gelatin, BSA and Tween 20 (0.2% each), in order to prevent non-specific
30 antibody staining. The cells or sections were then covered in Moviol 1-88 (Hoechst) and examined with a Zeiss Axiophot fluorescence microscopy. The photographs were prepared either conventionally or by computer-aided methods using a Kaf 1400 CCD camera (Photometric) and the
35 Adobe Photoshop 3.0 picture developing program.

In order to detect cytokeratin, vimentin and TGF β in human tumour tissue serial sections of frozen material were analysed immunohistochemically by the ABC method. The immunohistochemical analysis was carried out as described by Heider et al., 1995. Anti-cytokeratin (clone MNF 116; DAKO, Denmark), anti-vimentin (clone V9; DAKO, Denmark) and a mixture of anti-TGF β 1 and TGF β 2 (Santa Cruz, California) were used as the primary antibodies. The results of the staining were evaluated with a Zeiss Axioskop microscope.

For the simultaneous detection of vimentin and cytokeratin in tissue sections, double fluorescence analysis was carried out. Anti-vimentin (clone V9; DAKO, Denmark) and an anti-cytokeratin rabbit serum were used as primary antibodies, whilst Cy3-coupled anti-mouse IgG or FITC-coupled anti-rabbit IgG antibody were used as secondary antibodies. The fluorescence was evaluated using a Zeiss Axiophot 2 microscope with the aid of a Leica Quantimed Q500 picture analysis system.

f) RNA *in situ* hybridisation

For the RNA *in situ* hybridisation the frozen sections were fixed and extracted as described by Oft et al., 1993. For this the sections were fixed in 4% paraformaldehyde in PBS, washed twice in PBS, prehybridised for 2 hours and hybridised overnight with the appropriate S³⁵-labelled riboprobe at 52°C in 50% formamide, 0.6 M NaCl. After washing under stringent conditions (T_m -20°C) the sections were immersed in Kodak NTB liquid emulsion and illuminated for 2 weeks. The slides with the sections were counter-stained with haematoxylin/eosin and analysed under light and dark field illumination using a Zeiss Axiophot microscopes.

In order to prepare the S^{35} -labelled riboprobes the hTGF β 1-cDNA (R&D) and the cDNA for neomycin phosphotransferase, excised from a suitable retroviral vector (Redmond et al., 1988), were cloned into the
5 T₃.T₇ expression plasmids (Bluescript II KS Stratagene) and transcribed *in vitro*, in the presence of S^{35} -UTP for the antisense riboprobe and for the sense control probe.

The non-radioactive *in situ* hybridisation for TGF β in
10 sections of human tumour tissue was carried out by means of digoxigenin-labelled probes. The probe (hTGF β 1, see previous paragraph) was labelled by means of the DIG-RNA Labelling Kit made by Boehringer Mannheim according to the manufacturer's instructions. For the hybridisation frozen
15 sections (5-7 μ m) were fixed in 4% paraformaldehyde for 10 min, washed twice in PBS and subsequently acetylated for 10 min in 0.5 % acetic anhydride. After washing twice in PBS the sections were dehydrated in an ascending alcohol series, air-dried and subsequently incubated for 30 min at
20 52°C in a damp chamber. The hybridisation with the probe was carried out for 4 to 6 h at 52°C in a damp chamber. After hybridisation the slides were washed for 2 x 10 min in 2xSSC at 52°C and subsequently the bound probe was made visible by means of anti-digoxigenin antibody according to
25 the instructions of Boehringer Mannheim. The slides were briefly counter-stained with haematoxylin, covered and evaluated with a Zeiss Axioskop microscope.

g) Electron microscopy

30 The cells grown in the collagen gels were pre-fixed for 10 min in 3% paraformaldehyde in 0.2 M HEPES pH 7.3 at room temperature. The cells were further fixed on ice in 8% paraformaldehyde in 0.2 M HEPES pH 7.3 for 30-60 min.
35 For the immunocytochemistry the samples were dewatered in ethanol at ever lower temperatures, embedded in Lowicryl

HM20 or K4M and polymerised at -35°C by means of UV-light (Schwarz et al., 1993). To investigate the ultrastructure the cells were post-fixed with 1% osmium tetroxide in PBS pH 7.2 for 1 h on ice, stained with 1% aqueous uranyl acetate for 1 hour, dewatered in ethanol at room temperature and finally embedded in Epon. For the immunocytochemistry ultrathin sections were stuck to cover glasses (Schwarz, 1994). After the blocking of non-specific antibody binding sites with 0.5% bovine serum albumin and 0.2% gelatin in PBS the sections were incubated with rabbit-anti-Catenin-antibodies and subsequently with Cy3-labelled goat-anti-rabbit IgG. The labelled sections were stained with 4',6-diamino-2-phenylindole (DAPI), to make the nuclei visible under immunofluorescence microscopy.

h) Southern Blot analysis

Total DNA of cells or tumour material was isolated and processed using standard methods (Maniatis et al., 1982). DNA, extracted from cells before injection, from freshly excised tumour tissue (day 15 after the injection) and from tumour tissue which had been recultivated *in vitro* in the presence of G418 for 5 days, was digested with the restriction enzyme EcoRI (which cuts the retroviral vector only once), blotted onto a Gene Screen Membrane and hybridised with the cDNAs, coding either for the neomycin-phosphotransferase or for the v-Ha-ras gene.

i) Northern Blot analysis

The Northern Blot analysis was carried out as described by Chomczynski and Sacchi, 1987; as well as by Reichmann et al., 1992. Total RNA (10 µg per track) was placed on denaturing, formaldehyde-containing gels, blotted onto Gene Screen Membranes and hybridised with the entire

- 39 -

coding region of hTGF β 1-cDNA (R&D), which has sufficient homology with mTGF β 1, 2 and 3 to recognise all three mouse-TGF β -isoforms.

5 j) Semi-quantitative PCR

Total RNA from cells, grown on plastic, in collagen gels or from tumours, was isolated and processed for the semi-quantitative PCR. A TGF β 1-specific fragment was amplified
10 by means of RT-PCR under semi-quantitative conditions, using -actin primers as the internal control, as described by Leonard et al., 1993. For this the DNA was denatured at 94°C for 1 min, the primers were annealed at 65°C for 1 min, and the polymerase reactions were continued at 72°C
15 for 1 min. The amplification was continued for 20 and 30 cycles. The TGF β 1-specific primers TGGACCGCAA CAACGCCATC TATGAGAAAA CC (forward) and TGGAGCTGAA GCAATAGTTG GTATCCAGGG CT (reverse) (Clontech Inc.) were used. The results of the PCR were quantitatively evaluated on an
20 Image Quant Phospho-Imager. The values were standardised on the control product (-actin) and subsequently correlated with the value from control-3T3-fibroblast.

TGF β 1 was detected in tumour tissue by means of RT-PCR as
25 briefly described (Heider et al., 1996). The TGF β 1-specific oligonucleotide GCCCTGGACCACTATT GCTTC was used as 5'-primer and the TGF β 1-specific oligonucleotide TGCTCCACCTTGGGCTTGC was used as 3'-primer. The amplification products were separated on a 2 % ethidium
30 bromide-containing agarose gel and evaluated under UV-light by means of a video camera (MWG Biotech).

k) Transient transfection of a PAI-1-promoter-reporter gene construct.

35

A PAI-1 promoter-reporter construct (reporter gene was luciferase; 3TP-lux, Wrana et al., 1992) was transfected by lipofectamine transfection (Gibco) according to the manufacturer's instructions into CT26 cells or CT26-TBRII-dn cells. 8 hours after transfection TGF β 1 was added for 24 hours, while the controls were left without TGF β . Then cell lysates were prepared and the luciferase activity was measured in a Berthold Clinilumat as described by Wrana et al., 1992. In order to determine the transfection efficiency all the cells were cotransfected with an CMV- β -Gal reporter gene construct and the luciferase activities obtained were standardised by the β -Gal fluorescence intensity and by the protein concentration of the extract (Bradford).

15

1) Quantitative determination of soluble TGF β by means of ELISA assay

In order to determine the TGF β concentrations by means of ELISA assay, EpH4, polarised EpRas and fibroblastoid EpRas cells, isolated from tumours or converted *in vitro* with TGF β , were washed five times with PBS to remove exogenous TGF β and subsequently grown for 48 hours in serum-free DMEM. Then the cell culture supernatants were collected and the TGF β 1 concentrations were determined by means of a commercially obtainable ELISA-Kit (Promega; G1230) according to the manufacturer's instructions.

25

m) Immunoblots

In order to determine TGF β 1 in the tissue culture supernatants, 2 ml serum-free cell supernatants were concentrated by ultrafiltration (Centricon 10, Amicon) down to a final volume of 0.1 ml. The concentrated supernatants were mixed with 5-times concentrated SDS-PAGE probe buffer (without mercaptoethanol) and analysed by SDS-PAGE under non-reducing conditions. Equal aliquots of protein (50 μ g) were subjected to SDS-polyacrylamide gel electrophoresis; the immunoblot analysis was carried out as described by Hayman et al., 1993.

n) chicken embryo heart invasion assay

This assay was carried out as described by Behrens et al., 1993. In order to be able to distinguish invasive donor cells clearly from chicken heart cells, the test cells were charged with a vital fluorescent dye before examination. For this the cells were incubated for 1 hour in a glucose-containing Hanks saline solution containing 10 mM 5,6-carboxy-2',7'-dichlorofluorescein diacetate-succinimidyl ester (Molecular Probes) and 0.2×10^{-6} M Pluronic F127. In this way the fluorescent dye is covalently bound to intracellular proteins without affecting the viability or behaviour of the cells, determined by various differentiation and proliferation assays. The labelled cells were grown for 24 hours at high density, scraped off the plastic dish and brought into contact with pre-cultivated heart fragments of 9 day old chicken embryos on the surface of a soft agar layer. After 7 days' cultivation the fragments with the adhering cells were collected, flash-frozen in liquid nitrogen, frozen sections were prepared, fixed in methanol/acetone and the fluorescent cells were determined by epifluorescence microscopy (Axiophot, Zeiss).

o) Implantation of TGF β 1-charged Slow Release Pellets in mice

5 In order to expose Ras-transformed breast epithelial cells to activated TGF β 1 very early in tumour development, TGF β 1-charged Slow Release Elvax Pellets and either EpRas-epithelial cells or normal EpH4H cells were coinjected subcutaneously in mice. For control purposes pellets,
10 charged only with BSA, were coinjected. The pellets were prepared and charged according to the manufacturer's instructions.

15 Example 1

Ras expressing polarised epithelial cells undergo EF conversion during tumour development.

20 The tests carried out were suggested by the observation that Ras-transformed mouse-breast epithelial cells (EpRas cells) exhibit two completely different cell phenotypes. When they are grown on plastic substrates, these cells grow as ordered, dome-forming monolayers (hemicysts),
25 indicating a polarised epithelial phenotype (Fig. 1A, B). After being injected into mice, however, these same polarised cells formed tumours consisting of depolarised spindle-shaped cells with the capacity for invasive growth (Fig. 1A, C). In order to obtain further findings as to
30 the mechanisms underlying this phenotypical plasticity, the cell conversion observed was examined in detail by a combination of *in vivo* and *in vitro* experimental preparations. The cell clone EpH4 was used for this, which is derived from a well characterised mouse-breast
35 epithelial cell line (Reichmann et al., 1989; Reichmann et al., 1992; Strange et al., 1991). These cells exhibit a

stable polarised epithelial phenotype (Reichmann et al., 1994).

When suitable retroviral vectors were used tumorigenic subclones of EpH4 were formed by stable expression of the v-Ha-Ras-oncogene. After the expression of v-Ha-Ras as been confirmed by Western Blot analysis, cells from seven clones (referred to as EpRas-clones) were injected subcutaneously or directly into the mammary glands of Balb/c-mice. Tumours were formed regularly which were palpable 5-7 days after the injection of the cells.

The phenotype of these converted tumour cells was compared with that of the original differentiated clones: before the injection all seven EpRas-clones displayed the expected polarised phenotype (Fig. 1B and Table 1). By contrast, when cells were excised from the tumours and recultivated in the presence of G418, only converted, fibroblastoid cells were obtained (Fig. 1C, Table 1). Although they still expressed cytokeratin to a certain extent, these cells had lost many of their epithelial properties and acquired the expression of fibroblastic markers (Fig. 1C, Table 1). In order to demonstrate that the tumour cells came from the EpH4 donor cells originally injected, as well as to show that no rearrangement or reintegration of the Ras-containing retrovirus had taken place during the tumorigenesis and subsequent cultivation *in vitro*, the integration pattern of the retroviral constructs was determined by Southern Blot analysis. For this EpRas cells before injection, cells from a 15-day tumour and re-isolated cells from a 30-day tumour were analysed. When using probes with specificity for the neomycin resistance gene or the ras gene identical integration patterns were obtained in all three cell types (Fig. 1D).

- 44 -

The conversion of EpRas cells into fibroblastoid cells during tumour formation in mice is shown in Fig. 1.

Fig. 1A shows the principle of the strategy which was used in order to study EFC of Ras cells (7 different v-Ha-Ras expressing cell clones were used) *in vivo*.

Fig. 1B: Before injection cells of the clone Ep5 exhibited the formation of domes on plastic and staining both on E-cadherin (FITC, green fluorescence, appearing dark in the black and white representations), and also on cytokeratin (Texas-Red, red fluorescence). The common staining of both proteins on the periphery of the cell should be noted (yellow staining).

Fig. 1C: Ep5 cells, isolated from a tumour 28 days after the cell injection. These cells display a fibroblastoid appearance and express cytokeratin but no E-cadherin.

Fig. 1D: Southern Blot analysis. The EpRas-clone (Ep5), before injection (Ep5, plastic), removed from the tumour (Ep5, tumour), removed from the tumour and recultivated for 5 days in G418 (Ep5, ex tumour), shows the same retroviral integration pattern (detected with a neomycin-phosphotransferase(NPT) probe).

Example 2

Timing of EF conversion and behaviour of animal donor and receiver cells during tumorigenesis *in vivo*

Next, the stage of tumour development at which the subcutaneously injected epithelial EpRas cells undergo EF conversion, if at all, was examined. Three days after the injection the EpRas cells formed clearly defined nodules in which the cells expressed characteristic cytokeratins,

- 45 -

- but no vimentin (Fig. 2A). These epithelial cell nodules were already encapsulated by stroma cells (Fig. 2A). Cells which grew out of these microtumours on plastic and in the presence of G418, still demonstrated epithelial
- 5 properties. Seven days after the injection it was observed that the solid cell aggregation of Ep-Ras cells was beginning to break up at the edge of the tumour and the epithelial cells were mixed with vimentin-positive stroma cells at the periphery of the microtumours (by inward
- 10 migration of the stroma cells or outward migration of the donor cells). At this moment the donor cells still displayed epithelial properties, both in the tumour and also after isolation and *in vitro* cultivation in G418.
- 15 15 days after the injection three different cell types could be distinguished (Fig. 2C): about 20% of the tumour cells were green stained vimentin-positive stromal cells. Another 20% expressed only cytokeratins, indicating EpRas cells which have retained the epithelial phenotype. The
- 20 majority (50-60%) of the tumour mass, however, consisted of cells which co-expressed cytokeratin and vimentin. These cells are in all probability converted or converting EpRas cells. Both the epithelial and also the converted fibroblastoid cells were also obtained after G418
- 25 selection. Finally, the epithelial part could no longer be detected, either *in situ*, or on plastic, in five week old, fully developed tumours. By contrast parental EpH4 cells never formed tumours. When they were injected subcutaneously, the EpH4 cells developed into layers of
- 30 epithelial cells which sometimes formed lumina and cytokeratins, but expressed no vimentin (Fig. 2D). After a fairly long time these cells necrotised and were reabsorbed by the surrounding stroma.
- 35 In order to clearly identify the originally injected donor cells at the three different tumour stages, *in situ*

- 46 -

hybridisation was carried out on the neomycin resistance gene. These experiments showed that all cytokeratin expressing cells originated from donor cells. The frequency of donor cells relative to the stroma cells of the receiver animals increased with the size of the tumour and was greatest in fully developed tumours (Fig. 2E, F, G, H).

All in all, these data show that both the Ras expressing cells and also the epithelial control cells *in vivo* initially have an epithelial phenotype. As the development of the Ras cell tumours progresses the Ras-transformed cells progressively acquire fibroblastoid properties. By contrast the non-tumorigenic parental cells stably retain their epithelial properties until they die.

Fig. 2 shows the timing of the epithelial/mesenchymal conversion (EFC) during tumour development and demonstrates the fate of donor and receiver cells during this process.

Differently treated frozen sections of EpRas-tumours (clone Ep2) are shown, which were prepared on day 3 (Fig. 2A, E), on day 7 (Fig. 2B, F), on day 15 (Fig. 2C, G) and on day 28 (Fig. 2H) after the injection. The cell structures formed by non-tumorigenic EpH4 cells 15 days after the injection are shown in Fig. 2D. The sections were examined by immunofluorescence (Fig. 2A-D) and *in situ* hybridisation (Fig. 2E-H). The sections were double-stained with antibodies against a 46 kDa cytokeratin (Texas-Red, red fluorescence) and vimentin (FITC, green fluorescence). It was noted that in 3-day-old tumours the injected epithelial cells (stained red) and the mesenchymal cells of the host (stained green) are clearly separate. In 15-day-old tumours large numbers of cytokeratin/vimentin-double-positive cells are visible

- 47 -

(yellow-stained cells). These cells have undergone EFC. RNA-*in situ* hybridisation using a neomycin-phosphotransferase-probe confirms the donor origin of the tumour cells and shows the increasing density of the
5 tumour cells after EFC (Fig. 2E-H).

Example 3

TGF β 1 induces *in vitro* EF conversion in Ras expressing,
10 but not in normal epithelial cells

In order to identify the mechanism underlying the EF conversion, an experimental system was used with which EF conversion can be induced *in vitro* under defined and
15 physiologically relevant conditions. For this purpose normal EpH4 cells or Ras-transformed subclones of these cells (Ep-Ras clones) were grown in reconstituted collagen type I gels, using serum-free medium. These conditions made it possible to add defined polypeptide growth factors
20 and hormones which are known to be involved in the modulation of the epithelial phenotype. In gels of this kind, normal EpH4 cells developed into organ-like gland channels (tubuli) which frequently terminated in club-shaped hollow swellings. These structures looked very
25 similar to the end buds of the developing mammary gland which are formed by primary breast epithelial cells both in collagen gels and also *in vivo* (Fig. 3A). These structures could be induced to produce milk proteins efficiently by the addition of lactogenic hormones. When
30 these cells were isolated from the gel and grown on tissue culture plastic, they formed the expected regular epithelial monolayers, which formed recognisable domes, an indication that these cells are able to polarise efficiently (Fig. 3, right-hand Table).

35

Surprisingly, the EpRas-clones in these serum-free collagen gels also exhibited considerable lumen formation. The lumina were visible as early as 2-3 days after seeding. Thereafter more than 95% of these structures

5 developed relatively large cystic cavities (Fig. 3B, left-hand and centre Table) which resembled the alveoli of the fully developed milk-producing mammary gland. On plastic these cells in turn formed regular epithelial monolayers with domes, and thus exhibited the same epithelial

10 properties as the non-tumorigenic starting cells (Fig. 3B, right-hand Table).

The same EpRas cells behaved completely differently, however, when they were cultivated in 10% foetal calf

15 serum (FCS). Under these conditions they formed elongated, multi-cellular and invasively growing strings of cells which never showed any lumen formation. These strings consisted of non-polarised cells which had lost many epithelial properties (Fig. 3C and Fig. 4) and behave in a

20 strikingly similar manner to the ex vivo fibroblastoid tumour cells. These findings indicated that a factor contained in the FCS, co-operating with the activated Ha-Ras-oncoprotein, brings about the conversion of the epithelial EpRas cells into fibroblastoid cells.

25 In order to identify this factor or these factors, a number of growth factors (TGF β , heregulin, scatter-factor/hepatocyte growth factor, acidic and basic FGF, PDGF and TGF β 1) were added to the Ras-transformed cells

30 grown in collagen gels. Surprisingly, TGF β 1 was the only factor which showed striking and long-lasting effects on EpRas cells. When TGF β 1 was added, these cells grew into elongated, branching strings of cells similar to those induced by FCS. On tissue culture plastic these cells

35 exhibited a clear, fibroblastoid phenotype (Fig. 3D). In Eph4 control cells and other non-tumorigenic breast

epithelial-cell clones, by contrast, TGF β 1 was not able to induce EF conversion.

In order to examine whether the activity in the serum which promotes EF conversion is actually TGF β 1, cultures which contained 5% FCS were incubated with TGF β 1 neutralising antibodies. Under these conditions EpRas cells in turn formed cystic cavities very similar to those shown in Fig. 3B. Thus, the cell-converting activity present in FCS was identified as TGF β 1 and it was shown that TGF β 1 is the only or at least the predominant activity in FCS which can induce EF conversion.

Other ultrastructure and immunohistochemical analyses showed that most of the cystic structures consisted of a monolayer of polarised cells (Fig. 4A). These cells abundantly formed microvilli at their apical domain (the one facing the lumen), indicating a polarised organisation of the cells (Fig. 4A). Moreover, different types of epithelial-cell-typical cell-to-cell contact structures, i.e. tight junctions, characterised by the protein ZO-1, desmosomes (Fig. 4A) and the cell adhesion molecule E-cadherin typical of so-called "adherens junctions" (Fig. 4B) could be detected by their typical lateral or basolateral positions. Similarly, the protein β -catenin associated with E-cadherin showed basolateral localisation in most of the cells (Fig. 4C).

By contrast, the string-like cell structures induced by TGF β 1 consisted of loosely adhering spindle-shaped cells (Fig. 4D, inset picture). None of the epithelial marker proteins and ultrastructurally recognisable contact structures mentioned could be detected (Fig. 4D and Table 1), with the exception of a low, non-polarised expression of E-cadherin (Fig. 4E). The expression of β -catenin was greatly reduced and located chiefly in the

- 50 -

cytoplasm (Fig. 4F). Moreover, these cells expressed the expected mesenchymal markers (Table 1).

These results show that Ras-transformed mouse-breast epithelial cells exhibit exceptional plasticity in the phenotype, which ranges from epithelially polarised cells organised into ordered epithelia to fibroblastoid, migratory and invasively growing cells.

Fig. 3 shows the destruction of lumen formation and epithelial polarity by serum and TGF β 1.

Non-tumorigenic EpH4 cells (Fig. 3A) or tumorigenic EpRas cells (clone Ep5, Fig. 3B-D) were grown in collagen type I matrices. The macroscopically visible structures were photographed 8 days after plating out at low and high magnifications (left-hand and middle Table). Cells isolated from the gels and grown on tissue culture plastic are shown in the right-hand Tables.

Fig. 3A: Ep4H cells form channels and swellings resembling end-buds in serum-free collagen gels. On plastic these cells formed a regular epithelial monolayer and domes (hemicysts).

Fig. 3B: In serum-free collagen gels, wide channels and alveoli-like cysts are formed by EpRas cells.

Fig. 3C: Addition of 10% FCS causes the cells to form invasively growing irregular strings of cells without a lumen. On plastic these cells are similar to fibroblasts and are spindle-shaped.

Fig. 3D: TGF β 1 on its own (5 ng/ml) causes EpRas cells to grow into invasive strings of cells similar to those induced by FCS.

Fig.4 shows the breakdown of epithelial cell polarity in Ras-transformed breast epithelial cells after incubation with TGF β 1.

5

Alveoli-like cysts, formed by EpRas cells (clone Ep6) in serum-free collagen gels (Fig. 4A-C), and disordered strings of cells, formed by the same cells after treatment with TGF β 1 (Fig. 4D-F), were analysed for their epithelial organisation and formation of cell polarity. Sections through individual structures were photographed at high or low magnifications (inset pictures).

10

Fig. 4A: Transmission electron microscopy showed that the cysts obtained in the absence of TGF β 1 consisted of a monolayer of morphologically polarised cells which finally comprise the microvilli in their apical domain, facing the lumen (Fig. 4D). The inset picture shows a monolayer cyst of this kind at low magnification. By contrast, the strings of cells induced in the presence of TGF β 1 consist of loosely adhering cells without microvilli, desmosomes or tight junctions.

15

20

Fig. 4B, E: frozen sections through an alveolar cyst which were immunostained with an antibody against the cell adhesion molecule E-cadherin, showed clear basolateral localisation of the E-cadherins in most of the cells. In the TGF β 1-induced strings of cells, E-cadherin is reduced in its expression and is expressed over the entire surface of the fibroblastoid cells.

25

30

Fig. 4C, F: These show Lowicryl sections through structures similar to those shown in Fig. 4B and E, immunostained with an anti- β -catenin-antibody. The basolateral expression of β -catenin in most of the cells of the cyst (Fig. 4C) and the significantly reduced

35

β -catenin expression which is now localised predominantly in the cytoplasm should be noted (Fig. 4F).

Example 4

Fibroblastoid EpRas cells are invasive

EpRas cells which had undergone EFC showed signs of invasive behaviour in collagen gels. In order to obtain definitive proof of this invasive property, the chicken embryo heart invasion assays were used, the relevance of which to in vivo metastatisation has already been documented in detail (Mareel et al., 1979; Mareel, 1983). In this assay the migration of cells into embryo heart fragments was examined (Fig. 5A). In order to identify the penetrating cells clearly, they were labelled with a fluorescent dye (carboxy-dichloro-fluorescein-diacetate). During the incubation period of seven days no parental EpH4 cells migrated into the chicken heart tissue (Fig. 5A, B). In three different, fully-polarised Ep-Ras-clones, only a vanishingly small proportion of the cells were capable of migrating into the heart tissue (Fig. 5C). The few cells which migrated in were strongly stained with a vimentin antibody, but not with an anti-E-cadherin antibody. This confirms their conversion into a fibroblastoid phenotype, which is not surprising as the co-cultures contained serum. In contrast to the epithelial cells the fibroblastoid cells which had been obtained from tumours ("ex-Tu cells"), or cells which had been induced to EFC by the use of TGF β 1 in vitro, migrated into the heart muscle tissue in large numbers and relatively fast (Fig. 5D). These results show that EpRas cells are highly invasive after undergoing EFC, while non-converted epithelial cells exhibit only slight invasivity.

Fig. 5 shows the high invasivity of fibroblastoid EpRas cells in the chicken embryo heart invasion assay.

In vivo fluorescence-labelled cells were co-cultivated with chicken embryo heart fragments in order to test their invasivity, and sections through the fragments were examined histologically 7 days later. The non-tumorigenic epithelial starting cells (Eph4 cells) did not migrate into the heart fragments (Fig. 5A, B), and non-converted epithelial EpRas cells showed only slight invasivity (Fig. 5C). By contrast the converted, fibroblastoid cells obtained after TGF β 1 treatment were capable of migrating efficiently into the heart fragments (Fig. 5D).

Example 5

TGF β 1 maintains the fibroblastoid phenotype of converted EpRas cells through an autocrine loop

After it had been shown that TGF β 1 converts Ras-transformed epithelial cells into fibroblastoid cells, the question arose as to whether TGF β 1 is also involved in maintaining this phenotype. A possible explanation for the relative stability of the fibroblastoid phenotype (e.g. in cultures on plastic) was the autocrine production of fairly large amounts of TGF β 1 by the converted cells themselves. In order to settle this question, fibroblastoid EpRas cells were cultivated in extremely low concentrations in 1% FCS (in order to minimise the TGF β 1-concentration in the medium). Under these conditions individual cells grow into clearly spatially separated clones. As shown in Figures 6A and B, the clones obtained soon after plating out and consisting initially of a few cells had a fibroblastoid morphology at first. As the number of cell clones increased, the cells in the overwhelming majority of the clones gradually changed into

cells with an epithelial phenotype (Fig. 6A to D). This reversion was substantially complete 10 days after plating out (Fig. 6D).

- 5 In order to suppress the effects of autocrinally produced TGF β 1 completely and thereby definitively demonstrate that TGF β 1 is really necessary for maintaining the EF conversion, fibroblastoid cells (ex-tumour cells) isolated from a tumour were grown in the presence or absence of
- 10 TGF β 1 neutralising antibodies in collagen gels. In the absence of the antibodies the fibroblastoid tumour cells formed the expected thin, invasively growing strings of cells (Fig. 6E). The same cells, however, no longer grew invasively and developed into cystic structures consisting
- 15 of an epithelial monolayer when they were treated for eight days with the neutralising antibodies (Fig. 6F).

- Finally the amounts of TGF β 1-mRNA expressed in the cells and the TGF β 1-protein released into the culture medium
- 20 were determined. Three different EpRas clones as well as the parental Eph4 clone were grown for five days in collagen gels. When they were treated with 5 ng/ml TGF β 1, the EpRas-clones underwent EFC, while the similarly treated, non-transformed Eph4 cells retained their
- 25 epithelial phenotype. Analysis of these cells by semi-quantitative PCR (Fig. 7A) or by means of immunoblot (Fig. 7B) showed that the fibroblastoid cells induced by TGF β 1 produced quantities of TGF β 1-mRNA which were comparable with those of control fibroblasts (Fig. 7A).
- 30 This also applied to EpRas cells which had changed into fibroblastoid cells in tumours. By contrast, parental Eph4 cells and epithelial EpRas cells produced no or only small amounts of TGF β 1-mRNA (Fig. 7A). At the protein level essentially the same results were obtained when serum-free
- 35 culture supernatants were analysed by ELISA and Western blot (Fig. 7B).

Fig. 6 shows that TGF β 1 maintains the fibroblastoid phenotype of converted EpRas cells through an autocrine loop.

5

Fig. 6A-D: clones from fibroblastoid cells isolated from a tumour (ex-tumour cells) gradually change into clones consisting of epithelial cells. In order to produce the clones 500 cells per 100 mm dish were sown in medium containing 1% FCS. The medium was changed daily in order to dilute any autocrine factors. The same typical cell clone was photographed on day 1 (A), day 3 (B), day 5 (C) and day 10 (D) after plating out. The gradual transformation of the fibroblastoid cells into cells with an epithelial morphology is clearly visible.

Fig. 6E, F: fibroblastoid EpRas cells isolated from a tumour were selected for 5 days in G418 (in order to eliminate any cells originating from the receiver animal) and subsequently seeded into serum-free collagen gels. This was carried out either in the absence (E) or in the presence (F) of TGF β 1 neutralising antibodies. It can be seen that in the presence of a TGF β 1 neutralising antibody the tumour cells develop into lumen-shaped structures, whilst in the absence of the antibody they form the expected disordered strings of cells.

Fig. 7 shows that converted EpRas cells produce high concentrations of TGF β 1.

30

Fig. 7A: RNA from non-converted (epithelial) and converted (fibroblastoid) EpRas cells (clone Ep5) and also from non-tumorigenic Eph4 cells and NIH-3T3-fibroblasts (ATCC CRL 1658) was used for semi-quantitative PCR analysis. The significant increase in TGF β 1-mRNA in the fibroblastoid

35

cells should be noted. The TGF β 1 expression is recorded as a percentage of the values obtained with NIH-3T3 cells.

Fig. 7B: Similar results were obtained when the TGF β 1-concentrations in cell culture supernatants were analysed by Western Blot and ELISA (the numbers above the Western-Blot gel traces show the quantities of TGF β 1 in ng TGF β 1/ml determined in the ELISA). The data shown in Fig. 7B were confirmed with two other EpRas-clones (Ep2 and Ep6).

In all, these results indicate the major role of TGF β 1 not only in inducing EFC, but also in maintaining the fibroblastoid phenotype.

Example 6

Finally tests were carried out to determine whether TGF β 1 is actually expressed in EpRas tumours and whether TGF β 1 added experimentally in vivo can also bring about EFC and invasivity of the cells. Tumours growing from injected EpRas cells were examined for the expression of TGF β 1, 4 and 15 days after the injection of the cells, by RNA in situ hybridisation and immunohistochemistry. Just 4 days after the injection of the cells increased concentrations of TGF β 1-mRNA were detected at the outer edge of the nodes formed by the EpRas cells (Fig. 8A). The co-expression of TGF β 1 and neomycin phosphotransferase (NPT, which is expressed exclusively by the Ras-transformed donor cells) shown up by double immunofluorescence showed that the great majority of the donor cells (characterised by the red staining on NPT) produced no TGF β 1 (green staining) at this stage of the tumour development. On the other hand, cells of the surrounding tumour stroma originating from the receiver animal and non-epithelial in origin were distinctly positive for TGF β 1 (Fig. 8B). By contrast

tumours which had been removed 28 days after the injection showed a relatively high and uniform expression of TGF β 1-mRNA over the entire tumour region (Fig. 8C). In these tumours it was found that the injected EpRas cells

5 themselves produced TGF β 1 because they could be stained with antibodies against both NPT and TGF β 1; they displayed a yellow staining (Fig. 8D). Remarkably, most of the cells which produced TGF β 1 showed a reduced expression of cytokeratin, whereas the majority of the cells with high
10 cytokeratin expression could not be stained with antibodies against TGF β 1. This is further proof that the converted cells are actually those which also produce TGF β in the animal at advanced stages of the tumour.

15 These results show that host cells which surround the tumour tissue are able to initiate cell conversion. The converted tumour cells in turn themselves produce TGF β , thus speeding up cell conversion and subsequently the invasion processes.

20

In order to prove this directly, Slow Release Pellets charged with recombinant human TGF β 1 were applied close to the injected EpRas cells. The same TGF β 1 pellets, combined with non-tumorigenic Eph4 cells, were used as controls.

25 Surprisingly, EpRas cells located close to a TGF β 1 pellet were converted into irregularly shaped cells just 4 days after the injection and exhibited extensive migration into the surrounding host tissue. Surprisingly, even at this early stage, many of these cells were positive for
30 vimentin (Fig. 8F). By contrast, identical EpRas cells which had been injected in the absence of exogenous TGF β 1 formed smooth homogeneous nodes of vimentin-negative cells forming close cell contacts (Fig. 8E). As expected, TGF β 1 pellets located close to Eph4 cells could not noticeably
35 influence the phenotype of these non-tumorigenic cells. These *in vivo* data conform to the results obtained in

vitro and lead one to conclude that TGF β 1 has a key role in regulating the plasticity and invasivity of tumour cells.

- 5 Fig. 8 shows how TGF β 1 in experimentally induced tumours triggers the transition from the epithelial to the fibroblastoid state as well as the invasivity of the cells:
- 10 Fig. 8A-D: frozen sections of tumour stages on day 4 (A, B) and day 15 (C, D). The RNA *in situ* hybridisation shows that on day 4 the TGF β 1 expression is taking place in the outer periphery of the tumour (A), but on day 15 (C) it is occurring throughout the tumour. Arrow heads indicate the
- 15 boundary between tumour and surrounding stroma.

- Fig. 8B, D: frozen sections were stained with an anti-TGF β 1-antibody (green fluorescence) and an anti-neomycin phosphotransferase-antibody, which recognises the donor
- 20 cells (red fluorescence). The smaller diagrams show extracts at higher magnifications. It should be pointed out that at early stages of the tumour TGF β 1 is produced exclusively by the stroma around the tumour (B). By contrast, in 15 day-old tumours, TGF β is also expressed in
- 25 many donor cells within the tumour tissue (D, yellow fluorescence).

- Fig. 8E, F: Epithelial EpRas cells were injected subcutaneously into nude mice without (E) or together with
- 30 3-Elvax Slow Release Pellets charged with recombinant (active) TGF β 1 (F). Frozen sections obtained from 4 day old tumours were double-stained with antibodies against cytokeratin (red) and vimentin (green). What is noticeable is the dramatic migration of cells into the surrounding
- 35 tissue induced in the vicinity of the TGF β 1-releasing pellets (white circle).

Example 7

- Effect of TGF β 1 on normal breast epithelial cells: Control
5 of milk duct morphogenesis by regulating cell growth, cell
polarisation and apoptosis

Since the TGF β -super-family of polypeptide factors is
involved primarily in morphogenetic processes during
10 embryo development, the role of TGF β 1 in normal mammary
gland development was also examined within the scope of
the present invention. For this purpose normal breast
epithelial cells of the cell line EpH4 were sown in serum-
free collagen gels. Unlike in the experiments in Example
15 3, the serum needed during sowing for the collagen gel to
set and washed out one day later was specially selected
for a low content of TGF β 1. Under these conditions the in
vitro organogenesis was completely inhibited, and no
tubular structures were formed (Fig. 10 A). When low
20 concentrations of TGF β 1 (0.1 ng/ml) were added, the cells
were able to proliferate and form atypical structures
which generally lacked lumina (Fig. 10 B). Further
investigations showed, however, that these structures
expressed ZO-1, a tight-junction protein, on the inside.
25 Thus, these structures bore some resemblance to the end
buds of the developing mammary gland.

By contrast, higher concentrations of TGF β 1 (>0.25 ng/ml),
caused the normal epithelial cells to stop growing and die
30 off by programmed cell death (apoptosis) (Fig. 5C). This
is an important difference between the normal epithelial
cells and the Ha-Ras-containing cells. Whereas the latter
are not induced into apoptosis and undergo EFC without
exception even concentrations of TGF β 1 which are 20 times
35 higher (5 ng/ml), the TGF β 1 concentration which regulates
the morphogenetic processes in normal breast epithelial

cells is strictly laid down. Possibly, aberrant morphogenesis caused by excessively high TGF β 1 concentrations is prevented by the fact that growth inhibition and apoptosis are induced in the cells instead.

5

The fact that it was not possible to induce fully differentiated tubular structures consisting of polarised cells with low concentrations of TGF β 1, might be due to suboptimal culture conditions. On the other hand the complete organogenesis of tubular structures might depend on TGF β 1 only being present during certain phases of the organ development. In order to examine this, the cells were treated with 0.1 ng/ml TGF β 1, as described above, until structures had formed, then TGF β 1 was washed out of the collagen gel. Surprisingly, the atypical structures then reorganised themselves without lumina and formed well-shaped tubular structures with typical lumina (Fig. 10 D, transient TGF β 1). These results lead one to conclude (i) that TGF β 1 is absolutely necessary for in vitro organogenesis, (ii) that the concentration is critical, with higher concentrations leading to apoptosis, and (iii) that TGF β 1 only has to act on the cells during certain phases of the organ development. This normal function of TGF β 1 in the development of breast epithelial cells is completely changed in the Ras-transformed cells, with TGF β 1 here causing an extremely abnormal form of tissue reorganisation which causes a transition from the epithelial to the fibroblastoid state (EFC) over a wide range of concentrations.

30

The next step was then to look for indications that TGF β 1, analogously to these in vitro findings, also controls the morphogenesis and programmed cell death of mammary gland epithelia in vivo. For this, mammary glands in mice during puberty were subjected to histological analysis combined with in situ hybridisation using a probe against TGF β 1.

35

During this phase the virginal mammary glands grow into the surrounding fatty tissue (fat pad). Growth, differentiation and morphogenesis of the mammary gland produced start from a structure which is termed the end bud and contains undifferentiated, not yet fully polarised epithelial cells. Sections through the end bud (where the proliferation and subsequent organogenesis, such as e.g. the branching off of the milk ducts, take place) were compared with sections through fully differentiated ducts of the gland (cf. the schematic diagram in Fig. 11, centre). Whereas TGF β 1 is produced in the mesenchymal stroma which surrounds the growing end buds (Fig. 11, left-hand Table), no such production of TGF β 1 was seen in the stroma cells surrounding an already differentiated gland duct (right-hand Table). These findings largely correspond to the in vitro data, in which temporary, pulsed treatment of the breast epithelial cells with TGF β 1 was necessary for the tubular morphogenesis.

Similarly, there was an indication that TGF β also regulates the programmed cell death (apoptosis) of breast epithelial cells in vivo. During reversion of the mammary gland after ablactation the alveolar cells undergo mass apoptosis, whereas the cells in the gland ducts survive and are retained. New growth of the mammary glands during another pregnancy starts from these cells. In order to examine the possible involvement of TGF β 1 in this process, frozen sections through the dying alveolar zone of a mammary gland and through an adjacent gland duct region were prepared three days after the end of lactation (Fig. 12, representation in the centre). In the region which had just undergone apoptosis, the mesenchymal cells surrounding the dying alveoli expressed high concentrations of TGF β 1 (Fig. 12, left-hand Table), while the mesenchymal cells surrounding the surviving ductal structures expressed no TGF β 1 (Fig. 12, right-hand

Tables). In both cases it is probable that crosstalk takes place between the epithelial cells and the TGF β 1 production induced in the mesenchyme.

- 5 Fig. 10 shows that a low concentration of TGF β 1 controls the in vitro morphogenesis of normal mammary gland epithelial cells, particularly when the factor is given transiently. Higher concentrations of TGF β 1 cause apoptosis in the same cells.

10

- Normal EpRas cells were sown in collagen gels, using a foetal calf serum selected for a particularly low TGF β 1 content during the sowing. Under these conditions the cells do not form any tubular structures (Fig. 10A). In the presence of 0.1 ng/ml of TGF β 1 the cells form branched structures, but these lack lumina (Fig. 10B). If the TGF β 1 is removed from cultures with such structures on day 7 by washing, the cells form distinct hollow structures (Fig. 10D). Higher concentrations of TGF β 1 cause cell death (apoptosis, Fig. 10C, lower magnification on the left, higher magnification on the right).

20

- Fig. 11: shows the in vivo expression of TGF β 1 during the formation of the normal mammary gland during puberty (day 25)

25

- Frozen sections through end buds of a virginal mammary gland (left-hand panels) or through already formed gland ducts (right-hand panels) were prepared as shown in the central diagram. Successive sections in a series of sections were subjected to RNA in situ hybridisation for TGF β 1 mRNA (upper panels) or histologically stained. It is clearly apparent that mesenchymal cells which surround the end bud strongly express TGF β 1 (left-hand panels), whereas in cells which surround the differentiated gland ducts, there is no detectable TGF β 1 expression.

35

Fig. 12 shows the in vivo expression of TGF β 1 in the breakdown of the fully developed mammary gland after ablactation.

5

Young mice were taken away from their nursing mothers, thus triggering the reversion of the fully developed mammary gland. 3 days later frozen sections were taken through the dying areas of the mammary gland (left-hand panels) as well as through the gland ducts unaffected by the apoptosis (right-hand panels) (cf. the diagram in the centre of the Figure) The sections were then examined for TGF β 1 expression, as described in the legend to Fig. 11. Whereas TGF β 1 producing cells are clearly detectable in the area surrounding the dying alveoli (left-hand panels) there are none around the surviving gland ducts.

Example 8

20 Coexpression of vimentin and cytokeratins in human tumour tissue. Expression of TGF β by human primary tumours

In the previous Examples a well characterised cell model, namely Ras-transformed breast epithelial cells from the mouse, was used. It was thus very important to assess how far the results obtained with this model system as to the activity of the TGF β receptor on the phenotypical plasticity and invasivity of epithelial tumour cells applies to human carcinomas. For this purpose 31 kidney cell carcinomas and 64 breast tumours of different degrees of malignancy were examined by immunohistochemistry. Firstly, corresponding histological sections through such tumours were double-labelled with antibodies against general epithelial cytokeratins and antibodies against the mesenchymal marker vimentin. Tumour cells which coexpress both markers have probably undergone EPC. Secondly,

adjacent sections from the breast tumours were labelled with antibodies against human TGF β 1 and TGF β 2, in order to find out whether the tumour cells also produce TGF β .

- 5 As shown in the following Table, 74% of the kidney cell carcinomas expressed both cytokeratin and also vimentin in the degenerate epithelial tumour cells. As expected, the fibroblastoid cells of the tumour stroma expressed only vimentin, but no cytokeratin. In the breast carcinoma
10 cells the percentage of tumours which coexpressed cytokeratin and vimentin was smaller, namely between 24 and 27 %.

- The results of the histochemical analysis of the same
15 breast tumours for the expression of TGF β were even clearer. Here, all the tumours tested showed clear staining of the tumour cells for TGF β (Table). The tumour stroma was stained more weakly or not at all for most of the tumours. The specificity of the staining was also
20 clear from the staining of normal tissue; on the skin, for example, as expected, only the basal cell layers of the keratinocytes were positive. As is also shown in the Table, the results of the histochemical staining were also fully confirmed by in-situ hybridisation for TGF β as well
25 as by RT-PCR.

- These results show that for a significant proportion of the human tumours investigated, there were clear indications, in two ways, that the tumour cells
30 corresponding to the model in Fig. 9 had both undergone EFC and had also highly regulated the production of TGF β .

- The Table shows that human kidney cell and breast carcinomas coexpress cytokeratin and vimentin. This is a
35 clear indication that EFC has taken place. Similarly, all the tumours investigated produce TGF β .

- The upper part of Table (A) shows the results of the staining for cytokeratin and vimentin on frozen sections of the types of tumour specified. The lower part (B) shows the results of staining for TGF β on sections through the same breast tumours. Footnotes give the results of control experiments for the TGF β expression (by RT-PCR) and the expression of TGF β in the tumour stroma.

10

A Coexpression of vimentin and basal cytokeratins

Type of tumour	Subtype	Number of tumours analysed	Number of tumours with vim./cytok. coexpr.
kidney cell carcinomas (RCC)		31	23/31 (74%)
breast tumours			
	fibroadenoma (FA)	64	18/64 (28%)
	invasive ductal ca. (IDC)	3	2/3 (66%)
	invasive lobular ca. (ILC)	34	8/34 (24%)
	inv. ductal-lobular ca. (IDLC)	26	7/26 (27%)
		1	1/1

15

B Expression of TGF β -1/2

Type of tumour	Subtype	Antibody staining	In situ hybridisation
breast tumours			
	fibroadenoma (FA)	3/3 (100%)	
	invasive ductal ca. (IDC)	33/33 (100%)	13/13 (100%)
	invasive lobular ca. (ILC)	23/23 (100%)	9/9 (100%)
	inv. ductal-lobular ca. (IDLC)	1/1 (100%)	

- 66 -

Additional analyses

1. RT-PCR of IDC and ILC: 11 cases positive with Ab are also positive with RT-PCR
- 5 2. expression in the tumour stroma:
antibody: in 35/61 cases, slight staining
In situ hybridisation: in 3/22 cases

Example 9

10

Neutralising antibodies against TGF β prevent invasive growth of human tumour cell lines in collagen gel

- In Example 8 histochemical examination of sections through
- 15 tumour tissue provided evidence that the hypotheses reached with the model system regarding TGF β -induced EFC and the subsequent autocrine production of TGF β also apply to many human tumours. In order to obtain more direct
- 20 evidence of this, experiments were carried out to determine whether human tumour cells which grow invasively in the collagen gel can be converted into non-invasively growing cells by the administration of TGF β -neutralising
- 25 antibodies. The kidney carcinoma cell line MZ 1795 and the nasopharyngeal-carcinoma line KB were used. The cells of both lines grew in collagen gels containing 5% FCS without
- 30 TGF β -antibodies or after the addition of TGF β to form networks and strings of fibroblastoid cells (Fig. 13, right-hand panels). In the presence of TGF β neutralising antibodies (cf. Example 5, Fig. 6), on the other hand,
- the cells formed compact clumps without any reference to
- invasive growth (Fig.13, left-hand panels).

Fig. 13 shows that TGF β neutralising antibodies prevent the invasive growth of human tumour cell lines in collagen

35 gel.

MZ 1795 cells and KB cells were sown in serum-free collagen gels to which was added either 2% serum or 5 ng/ml of TGF β (+ TGF β , right-hand panels) or to which a

- 67 -

mixture of different antibodies against TGF β (- TGF β , left-hand panels; cf. Example 5, Fig. 6) was added. After 10 days microphotographs of the cells in the collagen gels were prepared. Whereas both the MZ 1795 cells (top panels, 5 higher magnification, bottom panels; summaries, lower magnification) and also the KB cells (lower panels) grow *en masse* into the collagen gel, when TGF β is present (right-hand panels), the same cells in the presence of TGF β -neutralising antibodies form compact clumps without 10 any cells growing out (left-hand panels).

Example 10

- 15 The expression of a dominant-negative TGF β receptor prevents EF conversion and slows down tumour growth of Ras-transformed breast epithelial cells

The most direct proof of the presumed mechanism of 20 activity (proof of principle) for the activity of TGF β receptor inhibitors in inhibiting tumour progression consists in demonstrating this activity directly in the tumour-bearing animal. This was not possible within the scope of these Examples with the TGF β -neutralising 25 antibodies used in vitro, as the large amounts of antibody needed for in vivo tests of this kind were not available. An alternative approach was therefore adopted.

There is a "kinase-dead" mutant of the human TGF β receptor 30 type II (T β RII-dn), which also acts as a dominant-negative receptor (i.e. one that switches off the function of the wild-type receptors). A cDNA of this T β RII-dn was expressed in Ras-transformed Eph4 cells (Ep-Ras) with the aid of retroviral vectors. The clones obtained grew very 35 slowly and required medium with a high (20%) serum content, in order to be capable of expansion. After

injection into nude mice these cells had formed either no tumours at all or only small tumours (Fig. 14, top) up to the time when the mice injected with control cells (Ep-Ras) had to be killed because of their excessively large tumours.

The tumour cells were isolated from the slowest-growing Ep-Ras-T β RII-dn tumour as well as from a control tumour induced by Ep-Ras cells and cultivated (cf. Example 1). Whilst the cells of the control tumour exhibited the expected fibroblastoid morphology (bottom of Fig. 14, left-hand Table), the tumour cells isolated from the slow-growing Ep-Ras-T β RII-dn exhibited a distinctly epitheloid morphology (bottom of Fig. 14, right-hand Table). This shows that the EF conversion occurring during tumour formation by Ep-Ras cells is inhibited by the expression of T β RII-dn and that this leads to a slowing down of tumour growth.

Fig. 14 shows that T β RII-dn expressing, Ras-transformed breast epithelial cells (Ep-Ras-T β RII-dn) exhibit slower growth in the animal and the cells isolated from these tumours have not undergone any EF conversion.

Four different clones of Ep-Ras-T β RII-dn cells as well as an Ep-Ras control clone were each injected subcutaneously into 3 nude mice (1×10^6 cells per animal). After 3 weeks the tumours were excised and weighed. The diagram in the upper part of Fig. 14 gives the mean values of the tumour weights obtained. The tumour cells from an Ep-Ras-T β RII-dn tumour, obtained from the slowest tumour-forming Ep-Ras-T β RII-dn clone, and an Ep-Ras control tumour were cultured, selected in G418 (cf. Example 1) and photographed after 10 days under phase contrast. The bottom left-hand panel shows the fibroblastoid cells which have grown from the Ep-Ras tumour, whereas the right-hand

panel shows the epitheloid cells which have grown from the Ep-Ras-T β RII-dn tumour.

5 Example 11

Expression of T β RII-dn in fibroblastoid, highly metastasising colon carcinoma cells (CT26): inhibition of the invasive growth of these cells in vitro, delaying of
10 tumour formation and inhibition of the formation of lung metastases in mice

Once it was shown that the dominant-negative TGF β receptor (T β RII-dn) could prevent both the EF conversion of EpRas
15 cells and also dramatically slowed down the tumour growth of these cells, it was useful to examine the efficacy of this T β RII-dn in tumour cells which had already stably undergone EF conversion and were already highly metastatic. The mouse colon carcinoma cell line CT26, an
20 established mouse model for lung metastasis formation from a primary tumour (Brattain et al. 1980), was chosen. These cells were infected with an T β RII-dn-expressing retrovirus (cf. Example 10), T β RII-dn-expressing clones were selected and various clones were subjected to analysis in vitro and
25 in vivo.

Two types of T β RII-dn-expressing CT-26 clones (CT26-T β RII-dn) were obtained. The first type showed a distinctly epithelial, but still abnormal morphology on plastic and
30 expressed small amounts of the epithelial markers E-cadherin and ZO-1. The second type of clone, on the other hand, on plastic formed lawns of cells with epithelial morphology which even formed hemicysts (domes). As expected, this second type of clone showed high lateral
35 expression of the epithelial markers E-cadherin and ZO-1. Control CT26 cells which were infected with a retrovirus

without an insert exhibited the expected fibroblastoid morphology on plastic and no expression of epithelial markers. It was thus shown that T β RII-dn is capable of converting the fibroblastoid CT26 cells into epithelial cells *in vitro* and thus effecting FE conversion.

Next, representative CT26-T β RII-dn clones of both types as well as CT26 control cells are sown in collagen gels with 5% FCS. Fig. 16 shows that the control cells grow into the expected enormous strings and networks of fibroblastoid cells (Fig. 15, top half of the picture, left-hand panels). By contrast the CT26-T β RII-dn clones of type 1 formed compact clumps with only a few single cells growing out (Fig. 15, top half of the picture, middle panels) whereas the CT26-T β RII-dn clones of type 2 only grew into tiny, compact groups of cells (Fig. 15, top half of the picture, right-hand panels). This showed that T β RII-dn prevents the invasive growth of CT26 cells in the collagen gel.

Then the same cell types were tested by the chicken heart invasion assay (cf. Example 4, Fig. 5). Whereas control-CT26 cells, as expected, grew very invasively in this assay (Fig. 16, bottom half of the picture, left-hand panel), the CT26-T β RII-dn clones of types 1 and 2 in this test were only slightly invasive or not invasive at all (Fig. 15, bottom half of the picture, middle and right-hand Table.)

These experiments show that T β RII-dn causes an FE conversion of CT26 cells and totally inhibits their invasive growth in two assay systems. It was therefore of great interest to examine the behaviour of these cells in the animal. Therefore, CT26 control cells as well as 6 CT26-T β RII-dn-clones of types 1 and 2 were injected into nude mice. Whereas the control animals had to be killed

- 71 -

after 2-3 weeks on account of their excessively large tumours, the tumour growth in mice was delayed with CT26-T β R11-dn-clones of type 1 by about 3-4 weeks, whereas in mice with CT26-T β R11-dn-clones of type 2 it was delayed by 6-10 weeks or totally inhibited for 24 weeks (end of experiment) (3 animals, data not shown in the Fig.). These results show that T β R11-dn can also dramatically delay the growth of CT26-primary tumours in some cases.

Next, the ability of the CT26-T β R11-dn cells to colonise the lung from a primary tumour and form metastases was examined. As shown in Fig. 17 (diagram in bottom half), mice were injected with CT26 control cells (3 mice) or 7 different CT26-T β R11-dn-clones (type 1 and type 2, 3 mice per clone) and the growth of palpable tumours was awaited. After a certain tumour volume (4 cm³) had been reached the primary tumour was excised so that no tumour cells remained at the injection site. The mice thus treated were examined for lung metastases after their death.

All control animals (3 mice) bearing CT26-tumours died after 2-4 weeks of lung metastases (Fig. 16, diagram in top half of picture, dotted line). By contrast the formation of lung metastases could not be detected in any of the animals injected with CT26-T β R11-dn-clones even after 18 weeks (Fig. 16, diagram in top half of picture, black lines). 5 animals in which there was a local recurrence of the primary tumour were not included in the evaluation.

These data clearly show that T β R11-dn fully inhibits the metastatisation of CT26-primary tumours. Finally, the stage of metastatisation which is inhibited by T β R11-dn was checked. It is possible that only the migration of the CT26 cells out of the primary tumour into the blood vessels is inhibited. However, the settling of

the cells out of the circulation and in the lungs might also be affected. The latter is important because, for example, more tumour cells might enter the circulation of a person when a tumour was surgically removed. In order to
5 test this, different quantities of CT26 control cells and a number of CT26-T β RII-dn clones of type 2 were injected intravenously into mice (3 animals per cell type). The animals were then examined for lung metastases after death. Preliminary tests showed that a mere 500 CT26 cells
10 per animal are sufficient to form lung metastases in this way. Therefore 10 and 100 times the quantity of both cell types were injected. Fig. 17 shows that after 14 days (50,000 cells) and 28 days (5,000 cells) the CT26 control cells had formed lung metastases in all the animals. By
15 contrast, even after 40 days, all the animals injected with CT26-T β RII-dn clones were still alive and had not yet formed any lung metastases, as confirmed on individual mice killed at this stage. Thus T β RII-dn can also prevent CT26 cells already in the circulation from settling in the
20 lungs.

Fig. 15 shows that T β RII-dn inhibits both the invasive growth of CT26 cells in the collagen gel, and also suppresses the invasivity of the same cells in the chicken
25 heart invasivity test.

For the first test (collagen gel assay, top half of the picture) CT26 control cells (CT26, left-hand panels) and a CT26-T β RII-dn clone of type I (middle panels) and type 2
30 (right-hand panels) were sown in collagen gels with 5% serum and after 10 days microphotographs of the collagen gels were prepared. The gels were photographed at two different magnifications (lower magnification, top panels, higher magnification, lower panels). Whereas the CT26
35 control cells grew into large network-shaped and string-like structures, consisting of spindle-shaped,

- 73 -

fibroblastoid cells (left-hand panels), the CT26-T β RII-dn type 1 cells formed compact clumps of cells with very few cells growing into the gel (middle panels). The CT26-T β RII-dn clones of type 2 form only tiny compact cell groups without any ability to grow into the collagen gel (right-hand panels).

For the chicken heart invasion assay (bottom half of the picture) the test cells were charged with a fluorescent vital dye, brought into contact with chicken heart fragments and examined histologically after 7 days (cf. methods and Example 4). The control cells migrated efficiently into the chicken heart fragment (left-hand Table, light-coloured groups of cells and strings on the side of the boundary between test cells and chicken heart fragment indicated by a dotted line labelled H). By contrast type 1 clones migrated only slightly into the chicken heart tissue (middle panel, cf. the few light-coloured cells in the area marked H), while the CT26-T β RII-dn cells of type 2 did not grow invasively at all (all the light-coloured cells remained outside the chicken heart fragments H (dotted line)).

Fig. 16 shows that the expression of T β RII-dn in CT26 cells blocks their ability to form lung metastases from a primary tumour.

The progress of the experiments is shown diagrammatically in the bottom half of the picture. 7 different CT26-T β RII-dn clones (type 1 and 2, CT26+T β RII-dn) as well as CT26 control cells were used in the test. Syngenic Balb-C mice (3 per cell type) were injected with 1×10^6 cells per animal and the growth of tumours was awaited. After the primary tumours reached a size of 4 cm³ they were surgically removed and after they had died the mice were examined for lung metastases. The results are shown in the

5 diagram (top half of the picture). Whereas the 3 control animals died within 4 weeks of lung metastases (dotted line), all the animals treated with CT26-T β RII-dn cells were still alive and free from lung metastases after 18 months (black lines). In the case of the line ending after 14 weeks (top diagram) the primary tumour reached the critical size so late that 18 weeks had not passed by the time the test ended.

- 10 Fig. 18 shows that T β RII-dn in CT26 cells also inhibits their ability to settle in the lungs from the bloodstream and form metastases there.

15 The diagram in the top part of the Figure shows the progress of the experiment. Syngenic Balb-C mice (3 per cell type and cell quantity) were injected intravenously (into the caudal vein) with CT26 control cells and several CT26-T β RII-dn clones. The Figure shows that all three mice treated with 5,000 or 50,000 control cells (CT26) had died
20 of lung metastases after 28 or 14 days (+), whereas all the animals injected with CT26-T β RII-dn clones were still alive free from lung metastases after 40 days (-).

25 Example 12

The activated TGF β receptor activates the transient transcription of a PAI-1- promoter-reporter gene construct, a process which is inhibited by T β RII-dn

- 30 With a view to discovering TGF β -(receptor) inhibitors by means of cellular assay in a High Throughput Screening (HTS) process a test cell is prepared as follows: a PAI-1- promoter-reporter gene construct is stably expressed in a
35 suitable cell (Ep-Ras or CT26). At the same time the human TGF β receptor chain (e.g.T β RII) selected for the screening

is expressed in this cell. By contrast, an unrelated receptor which also induces PAI-1 transcription, e.g. an FGF receptor, is expressed in control cells in addition to the PAI-1-reporter construct.

5 A prerequisite for the development of a test cell line of this kind is that the TGF β -induced PAI-I expression (suppressed by the inhibitor) is correlated with the tumour formation or metastasisation of the corresponding
10 cells in transient transfection tests. In order to check this, CT26 control cells and 5 CT26-T β RII-dn clones already tested in mice (cf. Example 11, Fig. 16) were transfected with the 3TP-lux PAI-1-reporter gene construct (Wrana et al., 1992), stimulated with TGF β or left
15 untreated and tested for PAI-1 expression (measurement of luciferase activity). As positive and negative controls a constitutively active TGF β R chain (T β RI (T204D); Wrana et al. 1994) as well as T β RII-dn DNA together with the PAI-1 reporter construct were co-transfected into untreated CT26
20 cells. Fig. 18 shows that untreated CT26 cells (CT26 controls) without TGF β treatment have a basal activity which corresponds to that of the negative controls (cotransfection of T β RII-dn). In the same control cells TGF β activates the reporter gene transcription to levels
25 which are attained in the positive controls by cotransfection of a constitutively active TGF β receptor.

Different T β RII-dn expressing CT26 clones behaved differently in this test (Fig.18, CT26-T β RII-dn 1-5). In
30 two clones (CT26-T β RII-dn 3 and 4) the luciferase activity achieved after TGF β stimulation was below or at the levels found in the negative controls, irrespective of whether the cells were tested before injection or after isolation from the very slow-growing tumour. In the three remaining
35 clones (in which the tumours induced grew faster) the luciferase activity was at the level of the negative

controls only before injection into the animal, after isolation from the tumour intermediate levels or even activities comparable to those in the positive controls were found (Fig. 18). It can be assumed that in the latter clones there was selection for cells in which the expression of T β RII-dn was down-regulated. This assumption was supported by the fact that renewed selection of the cells from the tumour in puromycin again killed off many cells and the surviving puromycin-resistant cells no longer exhibited increased PAI-1 transcription after TGF β stimulation. The results of these experiments show that the tumour formation/metastatisation in the animal is clearly correlated with the TGF β -activatability of a PAI-1 promoter-reporter gene construct.

Fig. 18 shows that the expression of T β RII-dn in CT26 cells suppresses the TGF β -induced transcription of a PAI-1 promoter-reporter gene construct.

CT26-control cells (CT26 controls) and 5 clones of CT26-T β RII-dn cells (CT26T β RII-dn 1-5) were transfected with a PAI-1 promoter-reporter gene construct (3TP-lux), the cells were stimulated with TGF β (+ TGF β) or left unstimulated (- TGF β) and the luciferase activity was measured in cell extracts. As positive controls the cDNA of a constitutively active TGF β receptor chain 1 (T β RI (T204D; Wrana et al., 1994) as well as the T β RII-dn-cDNA together with 3TP-lux were cotransfected into the cells. This measurement was carried out in cells before injection into the animal (before tumour induction) and after isolation and cultivation of the tumour cells for 3 days (isolated tumour cells) (cf. legend, box at top right). The bars indicate the standardised luciferase activity from extracts with the same protein content (cf. methods).

Bibliography

- Andrejauskas, E., and Moroni, C. (1989), *EMBO J.* 8, 2575-2581.
- 5 Antonelli-Orlidge, A., Saunders, K. B., Smith, S. R., and D'Amore, P. A. (1989), *Proc. Natl. Acad. Sci. U.S.A.* 86, 4544-4548.
- Aznavoorian, S., Murphy, A. N., Stetler, S. W., and Liotta, L. A. (1993), *Cancer* 71, 1368-1383.
- 10 Beham, A. et al., (1992), *Virchows Archiv A Pathol. Anat.*, Vol. 421, 209-215
- Behrens, J., Vakaet, L., Friis, R., Winterhager, E., Van, R. F., Mareel, M. M., and Birchmeier, W. (1993), *J. Cell Biol.* 120, 757-766.
- 15 Birchmeier, W., Weidner, K. M., and Behrens, J. (1993), *J. Cell. Sci. (Suppl)* 17, 159-164.
- Border, A. I., and Ruoslahti, V. (1992), *J. Clin. Invest.* 90, 1-7.
- 20 Bos, J. L. (1989), *Cancer Res.* 49, 4682-4689.
- Brattain, M.G., Strobels-Stevens, J., Find, D., Webb, M., and Sarraf, A. M. (1980), *Cancer Res.* 40, 2142-2452.
- Braunwalder et al. (1996), *Annal. Biochem.* 238, 159-164
- Buchmann, A., Ruggeri, B., Klein, S. A., and Balmain, A. (1991), *Cancer Res.* 51, 4097-4101.
- 25 Castle, V., Varani, J., Fligiel, S., Prochownik, E. V., and Dixit, V. (1991), *J. Clin Invest.* 87, 1883-1888.
- Caulin, C., Scholl, F. G., Frontelo, P., Gamallo, C., and Quintanilla, M. (1995), *Cell Growth & Differentiation* 6, 1027-1035.
- 30 Chomczynski, P., and Sacchi, N. (1987), *Anal. Biochem.* 162, 156-159.
- Clark, G. J., and Der, C. J. (1995), *Breast Cancer Research & Treatment* 35, 133-144.

- De Bortoli, M., Abou-Issa, H., Haley, B. E., and
Cho-Chung, Y. S. (1985), *Biochem. Biophys. Res.*
Commun. 127, 699-706.
- de Brito, P. A., Silverberg, S. G. and Orenstein, J. M.
5 (1993), *Hum. Pathol.* Feb. 24 (2), 132-42.
- Derynck, R., Jarret, J. A., Chen, E. Y., Eaton, D. H.,
Bell, J. R., Assoian, R. K., Roberts, A. B., Sporn, M.
B., and Goeddel, D. V. (1985), *Nature* 316, 701-705.
- Dvorak, H. F. (1986), *N. Engl. J. Med.* 315, 1650-1659.
- 10 Eaton, S., and Simons, K. (1995), *Cell* 82, 5-8.
- Edwards, D. R., Murphy, G., Reynolds, J. J., Whitham, S.
E., Docherty, A. J. P., Angel, P., and Heath, J. K.
(1987), *EMBO J.* 6, 1899-1904.
- Fearon, E. R., and Vogelstein, B. (1990), *Cell* 61,
15 759-767.
- Fakhrai H. et al. (1996), *Proc. Nat. Acad. Sci.* 93, 2909-
2914
- Fialka, I., Schwarz, H., Reichmann, E., Busslinger, M.,
and Beug, H. (1995), *J. Cell Biol.* 132, 1115-1132.
- 20 Frisch, S. M., and Francis, H. (1994), *J. Cell Biol.* 124,
619-626.
- Furth, M. E., Davies, L. J., Fleurdelys, B., and Scolnick,
E. M. (1982), *J. Virol.* 43, 294-304.
- Gabbert, H., Wagner, R., Moll, R., and Gerharz, C. D.
25 (1985), *Clin. exp. Metastasis* 3, 257-279.
- Guarino, M., Reale, D., Micoli, G. and Forloni, B. (1993),
Histopathology, May 22(5), 493-8.
- Guldborg, G. (1923), *Scand.* 4, 276-284.
- Hand, P., Thor, A., Wunderlich, D., Maurano, R., Caruso,
30 A., and Schlom, J. (1984), *Proc. Natl. Acad. Sci.*
U.S.A. 81, 5227-5231.
- Hartmann, G., Weidner, K. M., Schwarz, H., and Birchmeier,
W. (1994), *J. Biol. Chem.* 269, 21936-21939.
- Hayman, M. J., Meyer, S., Martin, F., Steinlein, P., and
35 Beug, H. (1993), *Cell* 74, 157-169.
- Heatly, M. et al., (1993), *J. Clin. Pathol.* 46, 441-445

- Heider, K.-H. et al. (1995), *Eur. J. Cancer*, 31A, 2385-2391
- Heider, K.-H. et al., 1996, *Virchows Archiv* 428, 267- 273
- Hoosein, N. M., McKnight, M. K., Levine, A. E., Mulder, K.
 5 M., Childress, K. E., Brattain, D. E., and Brattain,
 M. G. (1989), *Exp. Cell Res.* 181, 442-453.
- Jenkins, D. C., Stables, J. N., Wilkinson, J., Topley, P.,
 Holmes, L. S., Linstead, D. J., and Rapson, E. B.
 (1993), *Br. J. Cancer* 68, 856-861.
- 10 Kemler, R. (1993), *Trends Genet* 9:317-321.
- Kern, F. G., Cheville, A. L., and Liu, Y. L. (1990), *Semin
 Cancer Biol* 1, 317-328.
- Keski-Oja, J., Loef, E. B., Lyons, R. M., Coffey, R. J.
 J., and Moses, H. L. (1987), *J. Cell. Biochem.* 33,
 15 95-107.
- Kim, S. J., Jeang, K. T., Glick, A. B., Sporn, M. B., and
 Roberts, A. B. (1989), *J. Biol. Chem.* 264, 7041-7045.
- Kohl, N. E., Mosser, S. D., deSolms, S. J., Giuliani, E.
 A. et al., (1993), *Science* 260, 1934-1937
- 20 Kohl, N. E., Wilson, F. R., Mosser, S. D., Giuliani, E. et
 al., (1994), *Proc Natl Acad Sci USA* 91, 9141-9145
- Kohl, N. E., Omer, C. A., Conner, M. W., Anthony, N. J. et
 al., (1995), *Nature Med* 1, 792-797
- Kraus, M. H., Yuasa, Y., and Aaronson, S. A. (1984), *Proc.
 25 Natl. Acad. Sci. U.S.A.* 81, 5384-5388.
- Laiho, M., Saksela, O., and Keski-Oja, J. (1987), *J. Biol.
 Chem.* 262, 17467-17474.
- Land, H., Parada, L. F., and Weinberg, R. A. (1983),
Nature 304, 596-602.
- 30 LeJeune, S., Leek, R., Horak, E., Plowman, G., Greenall,
 M., and Harris, A. L. (1993), *Cancer Res.* 53, 3597-
 3602.
- Leonard, M. W., Lim, K. C., and Engel, J. D. (1993),
Development 119, 519-531.

- Lin H.Y.; Wang X.F.; Ng-Eaton E.; Weinberg R.A.; Lodish H.F. (1992) Cell 68(4), 775-785, erratum in Cell (1992) 70(6), 1068
- Liotta, L. A., Steeg, P. S., and Stetler-Stevenson, W. G. (1991), Cell 64, 327-336.
- Liotta, L. A., and Stetler-Stevenson, W. G. (1991), Cancer Res. 51, 5054-5059.
- Macauley, A., and Pawson, T. (1988), J. Virol. 62, 4712-4721.
- 10 Maniatis, T., Fritsch, E. F., and Sambrook, J. (1982), Molecular cloning. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- Mareel, M., Kint, J., and Meyvisch, C. (1979), Virchows Arch. (B) 30, 95-111.
- 15 Mareel, M. M. (1983), Cancer Metastasis Rev. 2, 201-218.
- Mareel, M. M. (1980), Int. Rev. exp. Pathology 22, 65-129.
- Mareel, M. M., Behrens, J., Birchmeier, W., De Bruyne, G. K., Vleminckx, K., Hoodgewus, A., Fiers, W. C., and Van Roy, F. M. (1991), Int. J. Cancer 47, 922-928.
- 20 Matsui et al., (1992), Hum. Pathol. Vol. 23, 1289-97
- Miettinen, P. J., Ebner, R., Lopez, A. R., and Derynck, R. (1994), J. Cell Biol. 127, 2021-2036.
- Monia, B.P. et al. (1996), Proc. Nat. Acad. Sci. 93, 15481-15484
- 25 Morgenstern, J., P. and Land, H. (1990), Nuc. Acid. Res. 18(12), 3587-3596
- Murthy, U., Anzano, M. A., and Creig, R. G. (1989), Int. J. Cancer 44, 110-115.
- Nishihara, K. and Tsuneyoshi, M. (1993), Hum. Pathol. Dec. 24(12), 1298-305.
- 30 Oft, M., Bohm, S., Wilczynski, S. P., and Iftner, T. (1993), Int. J. Cancer 53, 924-931.
- Parker, T. G., Packer, S. E., and Schneider, M. D. (1990), J Clin Invest 85, 507-514.
- 35 Payne, A., Olson, E. N., Hsiau, P., Roberts, R., Perryman, M. B., and Schneider, M. D. (1987), Proc. Natl. Acad. Sci. USA 84, 8956-8960.

- Pear, W. S., Nolan, G. P., Scott, M. L., Baltimore, D.
(1993) Proc. Natl. Acad. Sci. USA 90(18) 8392-8396
- Powell, S. M., Petersen, G. M., Krush, A. J., Booker, S.,
Jen, J., Giardiello, F. M., Hamilton, S. R.,
5 Vogelstein, B., and Kinzler, K. W. (1993), N. Engl. J.
Med. 329, 1982-1987.
- Quilliam, L. A., Kato, K., Rabun, K. M., Hisaka, M. M.,
Huff, S. Y., Campbell-Burk, S., and Der, C. J. (1994),
Mol Cell Biol 14, 1113-1121.
- 10 Redmond, S. M., Reichmann, E., Mueller, R. G., Friis, R.
R., Groner, B., and Hynes, N. E. (1988), Oncogene 2,
259-265.
- Reichmann, E. (1994), Semin Cancer Biol 5, 157-165.
- Reichmann, E., Ball, R., Groner, B., and Friis, R. R.
15 (1989), J. Cell Biol. 108, 1127-1138.
- Reichmann, E., Schwarz, H., Deiner, E. M., Leitner, I.,
Eilers, M., Berger, J., Busslinger, M., and Beug, H.
(1992), Cell 71, 1103-1116.
- Remington's Pharmaceutical Sciences, 1980, Mack Publ. Co.,
20 Easton, PA, Osol (ed.).
- Roberts, A. B., and Sporn, M. B. (1992), Cancer Surv. 14,
205-220.
- Roberts, A. B., Sporn, M. B., Assoian, R. K., Smith, J.
M., Roche, N. S., Wakefield, L. M., Heine, U. I.,
25 Liotta, L. A., Falanga, V., Kehrl, J. H., and Fauci,
A. S. (1986), Proc. Natl. Acad. Sci. U.S.A. 83, 4167-
4171.
- Sandford, K. K., Dunn, T. B., Westfall, B. B., Covallesky,
B. B., Covallesky, A. B., Dupree, L. T., and Earle, W.
30 R. (1961), J. Nat. Cancer Inst. 26, 1139-1193.
- Sato, Y., Tsuboi, R., Lyons, R., Moses, H., and Rifkin, D.
B. (1990), J. Cell Biol. 111, 757-763.
- Schoenenberger, C. A., Zuk, A., Zinkl, G. M., Kendall, D.,
and Matlin, K. S. (1994), J. Cell Sci. 107, 527-541.
- 35 Schultz-Cherry, S., Ribeiro, S., Gentry, L., and Murphy-
Ullrich, J. E. (1994), J Biol Chem 269, 26775-26782.

- Schwarz, H. (1994), Immunolabelling of ultrathin resin sections for immunofluorescence and electron microscopy. In *Electron microscopy 1994 ICEM*, B. Jouffrey and C. Colliex, eds. (Paris: Les editions de physique, Les Ulis, France), pp. 255-256.
- 5 Schwarz, H., Mueller-Schmidt, A., and Hoffmann, W. (1993), *Cell Tissue Res.* 273, 417-425.
- Seliger, B., Hühne, A., Knuth, A., Bernhard, H., Ehring, B., Tampé, R., and Huber, C. (1996), *Clin. Canc. Res.* 2, 1427-1433.
- 10 Slamon, D., J., de Kernion, J. B., Verma, I. M., and Cline, M. J. (1984), *Science* 224, 256-262.
- Sonnenberg, A., Daams, H., Calafat, J., and Hilgers, J. (1986), *Cancer Res.* 46, 5913-5922.
- 15 Soriano, J. V., Pepper, M. S., Nakamura, T., Orci, L., and Montesano, R. (1995), *J. Cell Sci.* 108, 413-430.
- Stacey, D. W., Feig, L. A., and Gibbs, J. B. (1991), *Mol. Cell. Biol.* 11, 4053-4064.
- Stoler, A. B., Stenback, F., and Balmain, A. (1993), *J. Cell. Biol.* 122, 1103-1117.
- 20 Strange, R., Li, F., Friis, R. R., Reichmann, E., Haenni, B., and Burri, P. H. (1991), *Cell. Growth Differ.* 2, 549-559.
- Thompson, A. M., Kerr, D. J., and Steel, C. M. (1991), *Br. J. Cancer* 63, 609-614.
- 25 Thompson, T. C., Truong, L. D., Timme, T. L., Kadmon, D., McCune, B. K., Flanders, K. C., Scardino, P. T., and Park, S. H. (1992), *J Cell Biochem Suppl.* 54-61.
- Thompson, T. C., Truong, L. D., Timme, T. L., Kadmon, D., McCune, B. K., Flanders, K. C., Scardino, P. T., and Park, S. H. (1993), *Cancer (Suppl.)* 71, 1165-1171.
- 30 Thorburn, A., Thorburn, J., Chen, S. Y., Powers, S., Shubeita, H. E., Feramisco, J. R., and Chien, K. R. (1993), *J. Biol. Chem.* 268, 2244-2249.
- 35 Vogelstein, B., and Kinzler, K. W. (1993), *Trends Genet.* 9, 138-141.

- Wargotz, E. S. and Norris, H. J. (1989), Cancer
Oct. 64(7), 1490-9.
- Wrana, J., L., Attisano, L., Cárcamo, J., Zentella, A.,
Doody, J., Laiho, M., Wang, X-F., and Massagué, J.
5 (1992) Cell, 71, 1003-1014
- Wrana, J., L., Attisano, L., Wieser, R., Ventura, F., and
Massagué, J. (1994), Nature 370, 341-347
- Welch, D. R., Fabra, A., and Nakajima, M. (1990), Proc.
Natl. Acad. Sci. U.S.A. 87, 7678-7682.
- 10 Whitman, M., and Melton, D. A. (1992), Nature 357,
252-254.
- Wong, S. Y., Purdie, A. T., and Han, P. (1992), Am. J
Pathol. 140, 1473-1482.
- Wright, J. H., McDonnell, S., Portella, G., Bowden, G. T.,
15 Balmain, A., and Matrisian, L. M. (1994), Mol.
Carcinog. 10, 207-215.
- Zambruno, G., Marchisio, P. C., Marconi, A.,
Vaschieri, C., Melchiori, A., Giannetti, A., and
De Luca, M. (1995), J. Cell Biol. 129, 853-865.